

# Jacobi Structures of Evolutionary Partial Differential Equations

Si-Qi Liu   Youjin Zhang

Department of Mathematical Sciences, Tsinghua University

Beijing 100084, P. R. China

Emails: liusq@mail.tsinghua.edu.cn, youjin@mail.tsinghua.edu.cn

March 29, 2011

## Abstract

In this paper we introduce the notion of infinite dimensional Jacobi structure to describe the geometrical structure of a class of nonlocal Hamiltonian systems which appear naturally when applying reciprocal transformations to Hamiltonian evolutionary PDEs. We prove that our class of infinite dimensional Jacobi structures is invariant under the action of reciprocal transformations that only change the spatial variable. The main technical tool is in a suitable generalization of the classical Schouten-Nijenhuis bracket to the space of the so called quasi-local multi-vectors, and a simple realization of this structure in the framework of supermanifolds. These constructions are used to compute the Lichnerowicz-Jacobi cohomologies and to prove a Darboux theorem for Jacobi structures with hydrodynamic leading terms. We also introduce the notion of bi-Jacobi structures, and consider the integrability of a system of evolutionary PDEs that possesses a bi-Jacobi structure.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Quasi-local multi-vectors</b>	<b>6</b>
2.1	The differential polynomial algebra $\mathcal{A}$ . . . . .	6
2.2	Differential operators on $\mathcal{A}$ . . . . .	12
2.3	Quasi-local multi-vectors . . . . .	14
2.4	The Schouten-Nijenhuis bracket . . . . .	18
2.5	Miura type transformations . . . . .	25
2.6	Reciprocal transformations . . . . .	29

<b>3</b>	<b>Jacobi structures</b>	<b>33</b>
3.1	Definition and examples . . . . .	33
3.2	Locality problem . . . . .	38
3.3	Deformation and cohomology . . . . .	40
3.4	Computation of cohomologies . . . . .	42
3.5	Bi-Jacobi structures . . . . .	45
<b>4</b>	<b>Examples</b>	<b>48</b>
4.1	The KdV equation v.s. the Camassa-Holm equation . . . . .	48
4.2	Bihamiltonian structures associated to Frobenius manifolds . . . . .	52
<b>5</b>	<b>Conclusion</b>	<b>54</b>

# 1 Introduction

The notion of Jacobi structure is a generalization of Poisson structures [37] on finite dimensional smooth manifolds, it has drawn much attention since its introduction by Lichnerowicz in 1978 [38]. Jacobi structure unifies several important mathematical structures which include Poisson manifolds, contact manifolds and locally conformal symplectic manifolds, it also gives a geometrical description of the local Lie algebra structures introduced by Kirillov [34]. On a manifold  $M^n$  endowed with a Jacobi structure (which is called a Jacobi manifold) one can define a bracket on the space of smooth functions  $C^\infty(M)$ , this bracket generalizes the usual Poisson bracket defined on a Poisson manifold, although it no longer satisfies the Leibniz rule, it still satisfies the Jacobi identity, and this is the main property that enable Jacobi structures to describe certain generalized Hamiltonian structures for dynamical systems. In this paper, we study a class of nonlocal Hamiltonian structures for evolutionary PDEs which have nice properties under reciprocal transformations. It turns out that this class of nonlocal Hamiltonian structures is the infinite dimensional counterpart of Jacobi structures on finite dimensional manifolds, so we call them Jacobi structures.

Hamiltonian structures for evolutionary PDEs are natural generalization of the ones for evolutionary ODEs, they have been playing important role in the study of qualitative properties of some nonlinear PDEs that arise in fluid mechanics and other fields of mathematics and physics [4]. They are also one of the main tools in the study of integrability of some physically important nonlinear evolutionary PDEs, see [12, 22, 28, 43, 49, 54] and references therein. The most frequently used class of Hamiltonian structures, often with an adjective “local”, for evolutionary PDEs with unknown functions  $u^1, \dots, u^n$  depending on the spatial variable  $x$  and time variable  $t$  are given by Poisson brackets of the form

$$\{F, G\} = \int \frac{\delta F}{\delta u^i(x)} P^{ij} \frac{\delta G}{\delta u^j(x)} dx, \quad (1.1)$$

where  $F, G$  are local functionals

$$F = \int f(u; u_x, \dots, u^{(m_1)}) dx, \quad G = \int g(u; u_x, \dots, u^{(m_2)}) dx$$

with densities  $f, g$  being differential polynomials of the unknown functions, i.e. they are polynomials in the  $x$ -derivatives of  $u^1, \dots, u^n$  with coefficients depending smoothly on  $u^i$ , and the operators  $P^{ij}, i, j = 1, \dots, n$  are polynomials in  $\partial_x = \frac{\partial}{\partial x}$  whose coefficients are differential polynomials. A Hamiltonian system of evolutionary PDEs associated to a local functional  $H$  have the representation

$$\frac{\partial u^i}{\partial t} = \{u^i(x), H\} = P^{ij} \frac{\delta H}{\delta u^j}, \quad i = 1, \dots, n.$$

By extending the notion of Schouten-Nijenhuis bracket defined on the space of multi-vectors of a finite dimensional manifold  $M$  to that of the infinite jet space  $J^\infty(M)$  of  $M$ , one can represent an infinite dimensional Hamiltonian structure of the above form in terms of a local Poisson bi-vector, and can define the associated Poisson cohomologies [21, 37, 49]. This leads to an analogue of the Darboux theorem on the normal forms of a class of infinite dimensional Poisson brackets which possess hydrodynamic limits [11, 21, 29]. Such normal forms play an important role in the study of the classification problem of bihamiltonian integrable hierarchies under Miura type transformations [16, 17, 21, 39], which are certain type of changes of the dependent variables of the hierarchies generalizing the well-known transformation between the Korteweg-de Vries (KdV) and the modified KdV equations [45].

Besides the *local* Hamiltonian structures, there exists an important class of *nonlocal* Hamiltonian structures which can also be represented by (1.1) with the operator  $P$  being replaced by an integro-differential operator, i.e. an operator that contains not only positive but also negative powers of  $\partial_x$ . For example, an operator  $P$  of the following form

$$P^{ij} = \sum_{s \geq 0} P_s^{ij} \partial_x^s + \sum_{\alpha, \beta=1}^{\ell} \eta^{\alpha\beta} X_\alpha^i \partial_x^{-1} X_\beta^j \quad (1.2)$$

defines a *weakly nonlocal Hamiltonian structure*, where  $P_s^{ij}, X_\alpha^i$  are differential polynomials, and  $(\eta^{\alpha\beta})$  is an  $\ell \times \ell$  constant symmetric matrix. Many important integrable PDEs, such as the KdV equation, the nonlinear Schrödinger equation and so on, have infinitely many weakly nonlocal Hamiltonian structures [44]. These nonlocal Hamiltonian structures have close relation to the integrability of these PDEs. Such nonlocal Hamiltonian structures also appear naturally when one applies certain reciprocal transformations to evolutionary PDEs which possess local Hamiltonian structures, see [9, 25] and references therein. Here the reciprocal transformations change the independent variables of the Hamiltonian systems, they play important role in the study of certain integrable systems such as the well-known Camassa-Holm equation [7, 8, 26, 27].

There is, however, a problem in the definition of the nonlocal Hamiltonian structures that needs to be clarified first: the action of the integral operator  $\partial_x^{-1}$  on the ring of

differential polynomials is not well-defined, so one can only use them in particular cases after adjusting the integration constants carefully. Our observation in the present paper is that, for an operator of the following special form

$$P^{ij} = \sum_{s \geq 0} P_s^{ij} \partial_x^s + X^i \partial_x^{-1} u_x^j + u_x^i \partial_x^{-1} X^j, \quad (1.3)$$

the corresponding weakly nonlocal Hamiltonian structure can be appropriately defined avoiding the use of the integral operator  $\partial_x^{-1}$ . In fact, if we introduce the following operator (which is called the *energy operator*, see Section 2.1)

$$E = \sum_{\alpha \geq 1} \sum_{s \geq 0} (-1)^s u^{i,\alpha} \partial_x^s \partial_{i,\alpha+s} - 1,$$

then we can show that

$$E \partial_x = 0, \quad \partial_x E = -u_x^i \frac{\delta}{\delta u^i}.$$

By using these identities, the Poisson bracket (1.1) defined by the operator (1.3) is reduced to the following form

$$\begin{aligned} \{F, G\}_P = & \int \frac{\delta F}{\delta u^i} \sum_{s \geq 0} P_s^{ij} \partial_x^s \frac{\delta G}{\delta u^j} \\ & + X^i E(F) \frac{\delta G}{\delta u^i} - \frac{\delta F}{\delta u^i} E(G) \quad dx. \end{aligned} \quad (1.4)$$

In this expression the operator  $\partial_x^{-1}$  does not appear, hence the above bracket is in fact a local object. The nonlocal Hamiltonian structures of the form (1.4) are called *quasi-local*, they are the main research object of the present paper.

The simplest quasi-local Hamiltonian structures are the ones with degree zero. It turns out that they coincide with Jacobi structures on the base manifold. For this reason, we name the quasi-local Hamiltonian structures Jacobi structures. When studying evolutionary PDEs, we are mainly interested in the Jacobi structures whose leading terms have degree one.

Jacobi structures of degree one have the following form:

$$P = g^{ij}(u) \partial_x + \Gamma_k^{ij}(u) u_x^k + V_k^i(u) u_x^k \partial_x^{-1} u_x^j + u_x^i \partial_x^{-1} V_k^j(u) u_x^k. \quad (1.5)$$

They have been investigated by Ferapontov [23] who proved that if  $P$  is a weakly nonlocal Hamiltonian structure and  $\det(g) \neq 0$ , then the inverse matrix of  $(g^{ij})$  defines a conformally flat metric  $(g_{ij})$ . In particular, when the dimension  $n \geq 3$ , this metric is flat if and only if the Jacobi structure  $P$  is a local Hamiltonian structure, i.e. the matrix  $(V_k^i)$  vanishes. By using Ferapontov and Pavlov's results [25], one can easily show that when  $n \geq 3$  there always exists a reciprocal transformation of the following form

$$d\tilde{x} = \rho dx + \sigma dt, \quad d\tilde{t} = dt \quad (1.6)$$

such that the transformed Jacobi structure is a local Hamiltonian structure, where  $\rho$  and  $\sigma$  are differential polynomials. We will prove that (see Section 3.1), when  $n = 1$  or  $2$ , such a reciprocal transformation also exists. Then according to the results of Dubrovin and Novikov [18, 19], the associated metric of a local Hamiltonian structure is flat, so there exists a set of flat coordinates such that the metric is given by a constant symmetric matrix  $(\eta_{ij})$ . The above discussion implies a classification theorem of Jacobi structures of degree one satisfying the condition  $\det(g) \neq 0$ , which are called Jacobi structures of hydrodynamic type.

**Theorem 1.1** *Let  $P$  be a Jacobi structure of hydrodynamic type, then there exists a reciprocal transformation  $\Phi$  and a Miura type transformation  $g$ , such that their composition transforms the Jacobi structure to  $g(\Phi(P)) = \eta^{ij}\partial_{\bar{x}}$ .*

We often denote  $P_{norm} = \eta^{ij}\partial_{\bar{x}}$ , and call it the normal form of the hydrodynamic Jacobi structure  $P$ . The main result of the present paper is a generalization of the above classification theorem to the full dispersive case.

**Main Theorem** *Let  $P$  be a Jacobi structure with hydrodynamic leading term, then there exists a reciprocal transformation  $\Phi$  and a Miura type transformation  $g$  such that  $g(\Phi(P)) = P_{norm}$ .*

The main technical tool for our study of the Jacobi structures is in a suitable generalization of the classical Schouten-Nijenhuis bracket to the space of the so called quasi-local multi-vectors, and a simple realization of this structure in the framework of supermanifolds, see Theorem 2.4.4. By using these constructions, we show in Theorem 3.3.6 that all the Lichnerowicz-Jacobi cohomologies of  $P_{norm}$  with positive degrees vanish, and the above Main Theorem is obtained as a corollary of this result. We note that when  $P$  is local, the Main Theorem also implies the Darboux theorem which was proved by Getzler and Degiovanni *et al* [11, 21, 29], our computation is much simpler than theirs.

By using the triviality of Lichnerowicz-Jacobi cohomologies, we also prove the following result concerning the relationship between bi-Jacobi structures (i.e. a pair of compatible Jacobi structures) and integrability of the associated evolutionary PDEs:

**Theorem 1.2** *If an evolutionary PDE possesses a bi-Jacobi structure with hydrodynamic leading term, then it has infinitely many conserved quantities and symmetries.*

This theorem is a reformulation of Lemma 3.5.2, it generalizes the similar result on the relationship between bihamiltonian structures and integrability.

In [16, 39], Dubrovin and the authors of the present paper studied the classification problem for semisimple bihamiltonian structures with hydrodynamic leading terms under the action of Miura type transformations. To any bihamiltonian structure of this type, we defined a set of functions of one variable which are called *central invariants*

(see Definition 3.5.4). These invariants, together with the leading terms of the bihamiltonian structure, determine the orbit of the bihamiltonian structure under the action of Miura type transformations. Now let us consider the transformation rule of the central invariants under the action of reciprocal transformations. By using the reciprocal transformation formula for Jacobi structures (see Theorem 2.6.3), we have the following result.

**Theorem 1.3** *If a reciprocal transformation transforms a local bi-Jacobi structure to a local one, then it preserves the central invariants of the local bi-Jacobi structure.*

This theorem is proved in Section 3.5, it has an interesting Corollary 3.5.7 which can be used to prove the equivalence of integrable systems that are obtained by applying different reciprocal transformations to a bihamiltonian integrable system, see Section 4.1 for illustrating examples.

We hope that the above mentioned results concerning the Jacobi structures together with some other results that will be given below could shed new light on the study of integrability of evolutionary PDEs and the geometry of infinite jet spaces.

The paper is organized as follows. In Section 2, we define the space of quasi-local multi-vectors and prove some useful formulae including the one for the Schouten-Nijenhuis bracket; then in Section 3 we study properties of Jacobi structures and bi-Jacobi structures. In Section 4, two examples are presented to illustrate the notion of Jacobi structure and their transformation rule under reciprocal transformations. We end the paper with some concluding remarks in Section 5.

## 2 Quasi-local multi-vectors

### 2.1 The differential polynomial algebra $\mathcal{A}$

Let  $M$  be a smooth manifold of dimension  $n$ ,  $J^\infty(M)$  be the infinite jet space of  $M$ . Recall that  $J^\infty(M)$  is a fiber bundle with fiber  $\mathbb{R}^\infty$  which is the projective limit of the projective system

$$\mathbb{R}^{kn} \underset{k \geq 1}{}, \quad \pi_{k,l} : \mathbb{R}^{kn} \rightarrow \mathbb{R}^{ln} \underset{k \geq l}{},$$

where the projection  $\pi_{k,l}$  is the forgetful map

$$\pi_{k,l} : (u^{(1)}, u^{(2)}, \dots, u^{(l)}, \dots, u^{(k)}) \mapsto (u^{(1)}, u^{(2)}, \dots, u^{(l)})$$

with  $u^{(k)} = (u^{1,k}, \dots, u^{n,k}) \in \mathbb{R}^n$ . Let  $U \times \mathbb{R}^\infty, V \times \mathbb{R}^\infty$  be two charts of a local trivializations of  $J^\infty(M)$  with local coordinates

$$U : (u^1, \dots, u^n; u^{(1)}, u^{(2)}, \dots), \quad V : (v^1, \dots, v^n; v^{(1)}, v^{(2)}, \dots),$$

then the transition functions of  $J^\infty(M)$  are given by the chain rule of higher order derivatives:

$$v^{i,1} = \frac{\partial v^i}{\partial u^j} u^{j,1}, \quad v^{i,2} = \frac{\partial v^i}{\partial u^j} u^{j,2} + \frac{\partial^2 v^i}{\partial u^k \partial u^l} u^{k,1} u^{l,1}, \quad \dots \quad (2.1)$$

We remark that the bundle  $J^\infty(M)$  is not a vector bundle, since the structure group is not the general linear group of the fiber.

A function  $f \in C^\infty(J^\infty(M))$  is called a differential polynomial if it depends on the jet variables polynomially in certain local coordinate system. All differential polynomials form a subalgebra of  $C^\infty(J^\infty(M))$ , we denote this subalgebra by  $\bar{\mathcal{A}}$ . Note that this definition does not depend on the choice of local coordinate system due to the form of the transition functions (2.1). On the ring  $\bar{\mathcal{A}}$  there is a gradation defined by

$$\deg u^{i,s} = s, \quad \deg f(u) = 0 \quad \text{for } f \in C^\infty(M). \quad (2.2)$$

We denote by  $\mathcal{A}$  the completion of  $\bar{\mathcal{A}}$  w.r.t. this gradation, and call it the differential polynomial ring of  $M$ . By abusing notations we also call elements of  $\mathcal{A}$  differential polynomials, though they may be infinite sums of differential polynomials (as defined above) with increasing degrees.

There is a vector field globally defined on  $J^\infty(M)$  which is defined as

$$\partial = \sum_{i=1}^n \sum_{s=0}^{\infty} u^{i,s+1} \frac{\partial}{\partial u^{i,s}}, \quad (2.3)$$

where  $u^{i,0} = u^i$ . This vector field yields a derivation on  $\mathcal{A}$  in an obvious way.

We assume henceforth that  $M$  is a contractible manifold with a local chart  $U : (u^1, \dots, u^n)$ , and all the computations are done on this chart. We denote  $\partial_{i,s} = \frac{\partial}{\partial u^{i,s}}$  if  $s \geq 0$ , and  $\partial_{i,s} = 0$  if  $s < 0$ .

In the remaining part of this subsection, we introduce several useful differential operators on  $\mathcal{A}$ , and prove some important properties of them. More details on the differential operators on  $\mathcal{A}$  can be found in the next subsection.

**Definition 2.1.1** For  $i = 1, \dots, n$  and integers  $\alpha, s \geq 0$ , the  $i$ -th higher generalized momentum operator of type  $(\alpha, s)$  is defined as

$$p_{i,\alpha,s} = \sum_{t \geq 0} (-1)^t \binom{t+s}{s} \partial^t \partial_{i,\alpha+s+t} : \mathcal{A} \rightarrow \mathcal{A}. \quad (2.4)$$

When  $s = -1$ , we denote  $p_{i,\alpha,-1} = \partial_{i,\alpha-1}$ .

When  $s = 0$ , the operator  $p_{i,\alpha,0}$  is just the generalized momentum operator in the Lagrangian mechanics on jet bundles. When  $\alpha = 0$ , the operator  $p_{i,0,s}$  coincides with the higher Euler operator  $\delta_{i,s}$  introduced by Kruskal, Miura, Gardner, and Zabusky [35] (see also [29]). In particular, if  $\alpha = s = 0$ , the operator  $p_{i,0,0}$  is just the  $i$ -th variational derivative  $\delta_i$ .

**Definition 2.1.2** For any integer  $s \geq -1$ , the higher energy operator is defined as

$$E_s = \sum_{\alpha \geq 1} u^{i,\alpha} p_{i,\alpha,s}. \quad (2.5)$$

In particular,  $E_{-1} = \partial$ . The energy operator is defined as  $E = E_0 - 1$ .

In Lagrangian mechanics, if we regard  $L \in \mathcal{A}$  as a Lagrangian, then  $E(L)$  is just the Hamiltonian of  $L$ , so we call  $E$  the *energy operator*.

**Lemma 2.1.3** *The operators  $p_{i,\alpha,s}$ ,  $E_s$ ,  $E$  satisfy the following identities:*

$$\begin{aligned}
& i) \quad p_{i,\alpha,s} \partial = p_{i,\alpha,s-1}, \quad E_s \partial = E_{s-1}, \quad E \partial = 0, \\
& ii) \quad \partial p_{i,\alpha,s} = p_{i,\alpha,s-1} - p_{i,\alpha-1,s}, \quad \partial E_s = E_{s-1} - u^{i,1} \delta_{i,s}, \quad \partial E = -u^{i,1} \delta_i, \\
& iii) \quad p_{i,\alpha,s}(f \cdot g) = \sum_{t \geq 0} (-1)^t \binom{t+s}{s} p_{i,\alpha,s+t}(f) \cdot \partial^t(g) + \partial^t(f) \cdot p_{i,\alpha,s+t}(g), \\
& \quad E_s(f \cdot g) = \sum_{t \geq 0} (-1)^t \binom{t+s}{s} E_{s+t}(f) \cdot \partial^t(g) + \partial^t(f) \cdot E_{s+t}(g), \\
& \quad E(f \cdot g) = \sum_{t \geq 0} (-1)^t E_t(f) \cdot \partial^t(g) + \partial^t(f) \cdot E_t(g) - f \cdot g, \\
& iv) \quad \sum_{t \geq 0} \binom{t+s}{s} \partial^t p_{i,\alpha,s+t} = \partial_{i,\alpha+s}, \\
& \quad \sum_{t \geq 0} \partial^t (f \cdot p_{i,\alpha,t}(g)) = \sum_{t \geq 0} \partial^t(f) \cdot \partial_{i,\alpha+t}(g), \\
& \quad \sum_{t \geq 0} \binom{s+t}{t} \partial^s E_{s+t} = \sum_{t \geq 0} \binom{\alpha+t}{t+1} u^{i,\alpha} \partial_{i,\alpha+t}, \\
& \quad \sum_{t \geq 0} \partial^t (f \cdot E_t(g)) = \sum_{t \geq 0} \sum_{\alpha \geq 1} \partial^t f u^{i,\alpha} \partial_{i,\alpha+t}(g).
\end{aligned}$$

*Proof* All these identities can be proved by direct computations. We omit the details here.  $\square$

Besides these identities, the operators  $p_{i,\alpha,s}$ ,  $E_s$ ,  $E$  also satisfy many other identities which are not easy to prove directly. The following lemma is quite useful when dealing with such identities.

**Lemma 2.1.4** *Let  $c : \mathcal{A} \rightarrow \mathcal{A}$  be a differential operator, if  $c(\mathcal{A}) \subset \mathbb{R}$  then  $c = 0$ . In particular, if  $\partial \cdot c = 0$ , then  $c = 0$ . Here we view  $\mathbb{R}$  as the subalgebra of  $\mathcal{A}$  consisting of constant functions.*

The proof of the lemma is a little bit technical, we give it in Section 2.2.

**Lemma 2.1.5** *The operators  $\delta_{i,t}$ ,  $\delta_j$ ,  $E_t$ ,  $E$  satisfy the following identities:*

$$i) \quad \delta_{i,t} \delta_j = (-1)^t \partial_{j,t} \delta_i, \quad (2.6)$$

$$ii) \quad E \delta_j = \partial_{j,0} E, \quad E_t \delta_j = (-1)^t \partial_{j,t} E \quad (t \geq 1), \quad (2.7)$$

$$iii) \quad E_t E = (-1)^t \sum_{s \geq 0} \binom{s+t}{t} \partial^s E_{s+t} E. \quad (2.8)$$

*Proof* The identity i) is just the closeness condition for a variational one-form. It can be proved by identifying the two sides with the following operator

$$(-1)^t \times_{p,q \geq 0} (-1)^p \binom{p}{t-q} \partial^{p+q-t} \partial_{i,p} \partial_{j,q}.$$

For the identity ii), note that

$$\partial E \delta_j = -u^{i,1} \delta_i \delta_j = -u^{i,1} \partial_{j,0} \delta_i = \partial \partial_{j,0} E,$$

so we have  $E \delta_j = \partial_{j,0} E$  according to Lemma 2.1.4. Next, for the  $t = 1$  case,

$$\partial E_1 \delta_j = E_0 - u^{i,1} \delta_{i,1} \delta_j = \partial_{j,0} E + \delta_j + u^{i,1} \partial_{j,1} \delta_i = \partial (-\partial_{j,1} E).$$

The  $t \geq 2$  cases can be proved similarly by induction on  $t$ .

The identity iii) can be proved as follow.

$$\begin{aligned} & (-1)^t E_t E = (-1)^t E_{t+1} \partial E = (-1)^{t+1} E_{t+1} u^{i,1} \delta_i \\ & = (-1)^{t+1} \times_{p \geq 0} (-1)^p \binom{p+t+1}{t+1} E_{p+t+1} u^{i,1} \partial^p \delta_i + u^{i,p+1} E_{p+t+1} \delta_i \\ & = (-1)^{t+1} \times_{p \geq 0} (-1)^p \binom{p+t+1}{t+1} u^{i,p+1} (-1)^{p+t+1} \partial_{i,p+t+1} E \\ & = \times_{\alpha \geq 1} \binom{\alpha+t}{t+1} u^{i,\alpha} \partial_{i,\alpha+t} E = \times_{s \geq 0} \binom{s+t}{t} \partial^s E_{s+t} E, \end{aligned}$$

here we used  $E_{p+t+1}(u^{i,1}) = 0$  and some identities given in Lemma 2.1.3.  $\square$

The following lemma is well-known in the theory of variational calculus, we give an alternative proof here in terms of notations we used above.

**Lemma 2.1.6** *Let  $f$  be a nonzero element of  $\mathcal{A}$ , then  $f \in \partial \mathcal{A}$  if and only if  $f \notin \mathbb{R}$  and  $\delta_i f = 0$  for all  $i = 1, \dots, n$ .*

*Proof* Since  $\delta_i \partial = 0$ , the necessary condition is obvious, we only need to prove the sufficient condition. Without loss of generality, we assume that  $f$  is a homogeneous element of  $\mathcal{A}$ . So we can set  $f = f(u, u^{(1)}, \dots, u^{(N)}) \notin \mathbb{R}$ . Introduce the notation  $Z_{(i,s)} = \partial_{i,s} Z$ . Then we have

$$(\delta_i f)_{(j,2N)} = (-1)^N f_{(i,N)(j,N)} = 0,$$

so  $f$  is linear in all  $u^{i,N}$  and we can assume that

$$f = \times_{i=1}^n f_i(u, u^{(1)}, \dots, u^{(N-1)}) u^{i,N} + f_0(u, u^{(1)}, \dots, u^{(N-1)}).$$

From the identity

$$(\delta_i f)_{(j, 2N-1)} = (-1)^N f_{(i, N)(j, N-1)} - f_{(i, N-1)(j, N)} = 0$$

it follows the existence of  $g(u, u^{(1)}, \dots, u^{(N-1)}) \in \mathcal{A}$  such that

$$f_i = g_{(i, N-1)},$$

thus  $f - \partial g$  is a differential polynomial that does not depend on  $u^{i, N}$ , then the proof of the lemma is finished by induction on  $N$ .  $\square$

We introduce the notation  $\sim$  as follows:

$$A \sim B, \text{ if } A - B \in \partial \mathcal{A},$$

where  $A, B \in \mathcal{A}$ . The following lemma is frequently used in this paper.

**Lemma 2.1.7** *Let  $X^i \in \mathcal{A}$  ( $i = 1, \dots, n$ ), if for any  $f \in \mathcal{A}$ , we have*

$$X^i \delta_i(f) \sim 0,$$

*then there exists  $c \in \mathbb{R}$  such that  $X^i = c u^{i, 1}$ .*

*Proof* Denote  $Z_j = \delta_j(X^i \delta_i f)$ , then  $Z_j = 0$ .

Without loss of generality, we assume that  $X^i$  ( $i = 1, \dots, n$ ) are homogeneous elements of  $\mathcal{A}$ . Let  $X^i = X^i(u, u^{(1)}, \dots, u^{(N)})$  which depends nontrivially on  $u^{(N)}$ . By taking  $f = u^i$  we see that  $X^i \sim 0$ , so  $X^i$  depends on  $u^{(N)}$  linearly and can be put into the form

$$X^i = \sum_{j=1}^N X_j^i(u, u^{(1)}, \dots, u^{(N-1)}) u^{j, N} + X_0^i(u, u^{(1)}, \dots, u^{(N-1)}).$$

If  $N = 2p$  is even, then by putting  $f = (-1)^p (u^{i, p})^2 / 2$  we get

$$Z_{i, (k, 2N)} = X_k^i + \delta^{ik} X_i^i = 0,$$

so  $X_j^i = 0$ , this contradicts our assumption on  $X^i$ .

Now let  $N$  be odd. We first assume that  $N = 2p + 1 \geq 3$ . By taking  $f = \frac{(-1)^{p+1}}{2} (u^{i, p+1})^2$  we obtain

$$Z_{i, (k, 2N+1)} = X_k^i - \delta^{ik} X_i^i = 0,$$

it follows that  $X_k^i = 0$  if  $k \neq i$ . Next, from

$$Z_{i, (i, 2N), (k, N)} = 1 + 2\delta^{ik} X_{i, (k, N-1)}^i = 0$$

we know that  $X_i^i$  does not depend on  $u^{(N-1)}$ . Then we have

$$Z_{i, (i, 2N)} = 2X_{0, (i, N-1)}^i + \partial X_i^i = 0.$$

Now let us take  $f = \frac{(-1)^{p+1}}{6} (u^{i,p+1})^3$ , then from

$$Z_{i,(i,2N)} = u^{i,p+1} 2X_{0,(i,N-1)}^i + \partial X_i^i - N X_i^i u^{i,p+2} = 0$$

it follows that  $X_i^i = 0$ . This contradicts our assumption that  $X^i$  depends nontrivially on  $u^{(N)}$ , so we must have  $N \leq 1$ .

If  $N = 1$ , similar to the case when  $N$  is odd and  $N \geq 3$ , we can obtain

$$X^i = c^i u^{i,1},$$

where  $c^i \in \mathbb{R}$ . Finally we take  $f = f(u)$ , then

$$Z_j = \prod_{i=1}^j (c^i - c^j) f_{(i,0)(j,0)} u^{i,1} = 0,$$

so  $c^i = c^j$ . The lemma is proved.

Finally, we remark that when  $X^i = c u^{i,1}$ ,

$$X^i \delta_i(f) = c u^{i,1} \delta_i(f) = \partial(-c E(f)) \sim 0,$$

so the condition of the lemma is indeed satisfied. □

The following two lemmas are very useful when computing  $\delta_i$  and  $E$ .

**Lemma 2.1.8** *Let  $D$  be a first order differential operator over  $\mathcal{A}$ , if*

$$D\partial = \partial(D - D(1)),$$

*then for any  $f \in \mathcal{A}$ , we have*

$$D(f) \sim X^i \delta_i(f) - a E(f),$$

*where  $a = D(1)$ ,  $X^i = (D - D(1))(u^i)$ .*

*Proof* The general form of  $D$  is

$$D = a + \prod_{i,s} X^{i,s} \partial_{i,s},$$

where  $X^{i,0} = X^i$ . The condition  $D\partial = \partial(D - D(1))$  is equivalent to

$$X^{i,s+1} = \partial X^{i,s} - a u^{i,s+1},$$

which implies

$$X^{i,s} = \partial^s X^i - \prod_{t=1}^s \partial^{s-t} a u^{i,t}.$$

Then by using the definition of  $\delta_i$  and  $E$ , the lemma is proved immediately. □

**Lemma 2.1.9** *Let  $f \in \mathcal{A}$  be a differential polynomial, if for any  $g \in \mathcal{A}$ ,  $fg \sim 0$ , then we have  $f = 0$ .*

*Proof* First we take  $g = 1$ , then there exists  $h \in \mathcal{A}$  such that  $f = \partial(h)$ . Next we take  $g = f$ , so we have  $\delta_i(f^2) = 0$ . Without loss of generality, we assume that  $h$  is a homogeneous differential polynomial, so there exists  $N \in \mathbb{N}$  such that  $h_{(i,s)} = 0$  for any  $s > N$ . On the other hand, we have

$$0 = \delta_i(f^2)_{(i,2N+2)} = (-1)^{N+1} f^2_{(i,N+1)(i,N+1)} = 2(-1)^{N+1} h_{(i,N)}^2,$$

so  $h$  must be a constant, hence  $f = 0$ . The lemma is proved.  $\square$

**Corollary 2.1.10** *Let  $f \in \mathcal{A}$  be a differential polynomial, and  $g_1, \dots, g_n, h \in \mathcal{A}$ . If for any first order differential operator  $D$  satisfying  $D\partial = \partial(D - D(1))$ , the following relation holds true*

$$D(f) \sim X^i g_i - a h,$$

where  $a = D(1)$ ,  $X^i = (D - D(1))(u^i)$ , then we have  $g_i = \delta_i(f)$  and  $h = E(f)$ .

## 2.2 Differential operators on $\mathcal{A}$

In this subsection, we define the algebra of differential operators over  $\mathcal{A}$ , and prove Lemma 2.1.4. The readers who are not interested in the proof of Lemma 2.1.4 can skip this subsection.

**Definition 2.2.1** *We define the following algebras*

$$\begin{aligned} \mathcal{D}_M &= C^\infty(M) \otimes \mathbb{R}[\partial_{i,0} | i = 1, \dots, n], \\ \mathcal{D}'_M &= \mathcal{A} \otimes \mathbb{R}[\partial_{i,0} | i = 1, \dots, n], \\ \bar{\mathcal{D}} &= \mathcal{A} \otimes \mathbb{R}[\partial_{i,s} | i = 1, \dots, n, s \geq 0], \end{aligned}$$

whose elements are called differential operators over  $M$ , differential operators over  $M$  with coefficients in  $\mathcal{A}$ , and bounded differential operators over  $\mathcal{A}$  respectively.

The algebra  $\bar{\mathcal{D}}$  possesses a gradation

$$\bar{\mathcal{D}} = \bigsqcup_{d \in \mathbb{Z}} \bar{\mathcal{D}}_d, \quad \bar{\mathcal{D}}_d = \left\{ \prod_{I,S} f^{I,S} \partial_{I,S} \mid \deg f^{I,S} - |S| = d \right\},$$

which induces a decreasing filtration

$$\bar{\mathcal{D}} \supset \cdots \supset \bar{\mathcal{D}}_{(d-1)} \supset \bar{\mathcal{D}}_{(d)} \supset \bar{\mathcal{D}}_{(d+1)} \supset \cdots, \quad \bar{\mathcal{D}}_{(d)} = \prod_{i \geq d} \bar{\mathcal{D}}_i,$$

we denote the topological completion of  $\bar{\mathcal{D}}$  w.r.t. the above filtration by  $\mathcal{D}$ , and call it the differential operator algebra over  $\mathcal{A}$ .

Here we used the multiple index notations:

$$(I, S) = (i_1, s_1; \dots; i_m, s_m), \quad \partial_{I,S} = \partial_{i_1, s_1} \cdots \partial_{i_m, s_m},$$

where  $m = 0, 1, 2, \dots$ ,  $|I| = m$ , and  $|S| = s_1 + \dots + s_m$ .

Before proving Lemma 2.1.4, we need some preparations.

**Lemma 2.2.2** *Let  $D \in \mathcal{D}_M$ , if for any  $f \in C^\infty(M)$ ,  $D(f) \in \mathbb{R}$ , then  $D = 0$ .*

*Proof* The general form of  $D$  reads

$$D = \sum_{|K| < m} a_K \partial^K,$$

where  $K = (k_1, \dots, k_n)$  is the multiple index, and

$$\partial^K = \partial_{1,0}^{k_1} \cdots \partial_{n,0}^{k_n}, \quad |K| = k_1 + \dots + k_n.$$

By choosing sufficiently many linearly independent  $f_1, \dots, f_N$  and by using the property of Wronskian determinants, one can prove that all the coefficients  $a_K$  vanish. The lemma is proved.  $\square$

**Lemma 2.2.3** *Let  $D \in \mathcal{D}'_M$ , if for any  $f \in C^\infty(M)$ ,  $D(f) \in \mathbb{R}$ , then  $D = 0$ .*

*Proof* Suppose

$$D = \sum_{I,S} u^{I,S} D_{I,S},$$

where  $(I, S) = (i_1, s_1; \dots; i_m, s_m)$  with  $s_i \geq 1$  and  $D_{I,S} \in \mathcal{D}_M$ , then every  $D_{I,S}$  satisfies the condition of Lemma 2.2.2, so  $D_{I,S} = 0$ , then  $D = 0$ . The lemma is proved.  $\square$

*Proof of Lemma 2.1.4* According to the definition of  $\mathcal{D}$ , every  $c \in \mathcal{D}$  can be decomposed into

$$c = \sum_{m \geq 0} c_m, \quad c_m = \sum_{|I|=m} \sum_S c_m^{I,S} \partial_{I,S},$$

where  $(I, S) = (i_1, s_1; \dots; i_m, s_m)$  with  $s_i \geq 1$ ,  $c_m^{I,S} \in \mathcal{D}'_M$ .

First we take  $f \in C^\infty(M)$ , so

$$c(f) = c_0(f),$$

since  $c(f) \in \mathbb{R}$ , according to Lemma 2.2.3, we have  $c_0 = 0$ .

Next, let  $m$  be a positive integer, assume that for any multiple index  $(I', S')$  with  $|I'| = m' < m$  we have proved  $c_{m'}^{I',S'} = 0$ , then for a multiple index  $(I, S)$  with  $|I| = m$ , let  $f = \tilde{f} u^{I,S}$  where  $\tilde{f} \in C^\infty(M)$ , then we have

$$c(f) = C(I, S) c_m^{I,S}(\tilde{f}),$$

where  $C(I, S) = \partial_{I,S}(u^{I,S})$  is a positive constant. Since  $c(f) \in \mathbb{R}$  for any  $\tilde{f}$ , according to Lemma 2.2.3, we have  $c_m^{I,S} = 0$ .

The lemma is proved by induction on  $m$ .  $\square$

## 2.3 Quasi-local multi-vectors

Let  $\mathcal{F}$  be the quotient space  $\mathcal{A}/\partial\mathcal{A}$  whose elements are called local functionals. The coset of  $f \in \mathcal{A}$  in  $\mathcal{F}$  is denoted by  $\int f dx$ , and  $f$  is called a density of this local functional. The space  $\mathcal{F}$  has a gradation induced from  $\mathcal{A}$  since  $\partial$  is homogeneous of degree one. Note that the operators  $\delta_i, E$  satisfy  $\delta_i\partial = 0, E\partial = 0$ , so they induce maps from  $\mathcal{F}$  to  $\mathcal{A}$  which we still denote by  $\delta_i$  and  $E$ .

We denote  $\mathcal{V}^p = \text{Alt}^p(\mathcal{F}, \mathcal{F})$  where  $\text{Alt}^p(V_1, V_2)$  stands for the linear space of all alternating  $p$ -linear maps from a linear space  $V_1$  to another linear space  $V_2$ . The elements of  $\mathcal{V}^p$  are called the generalized  $p$ -vectors over  $\mathcal{F}$ . We also use the notations  $\mathcal{V}^0 = \mathcal{F}, \mathcal{V}^{<0} = 0$ , and  $\mathcal{V} = \bigcup_{p \geq 0} \mathcal{V}^p$ . The quasi-local multi-vectors that we are going to define below are certain kind of generalized multi-vectors over  $\mathcal{F}$  which are generated by the maps  $E$  and  $\delta_i$ .

**Definition 2.3.1** *Let  $P \in \mathcal{V}^p$  be a generalized  $p$ -vector, we say that  $P$  is quasi-local if the action of  $P$  on  $F_1, \dots, F_p \in \mathcal{F}$  takes the following form*

$$\begin{aligned} P(F_1, \dots, F_p) = & \int Q_{s_1 \dots s_p}^{i_1 \dots i_p} \partial^{s_1} \delta_{i_1}(F_1) \dots \partial^{s_p} \delta_{i_p}(F_p) \\ & + R_{t_1 \dots t_{p-1}}^{j_1 \dots j_{p-1}} \bigotimes_{k=1}^p (-1)^{k-1} E(F_k) \partial^{t_1} \delta_{j_1}(F_1) \dots \hat{F}_k \dots \partial^{t_{p-1}} \delta_{j_{p-1}}(F_p) dx, \end{aligned}$$

where  $Q_{s_1 \dots s_p}^{i_1 \dots i_p}, R_{t_1 \dots t_{p-1}}^{j_1 \dots j_{p-1}} \in \mathcal{A}$ .  $P$  is called local if the second term of the above expression does not appear.

In order to give a more simple and concrete description of the class of quasi-local multi-vectors, we introduce a family of super variables  $\theta_i^s, \zeta$ , where  $i = 1, \dots, n, s = 0, 1, 2, \dots$ , and define

$$\mathcal{S} = \mathcal{A} \otimes \wedge^*(V), \text{ where } V = \bigoplus_{i,s} (\mathbb{R}\theta_i^s) \oplus \mathbb{R}\zeta.$$

The gradation on  $\mathcal{A}$  can be extended to a gradation on  $\mathcal{S}$  as follow:

$$\deg u^{i,s} = \deg \theta_i^s = s, \deg f(u) = \deg \theta_i = \deg \zeta = 0, \quad (2.9)$$

we still denote the completion of  $\mathcal{S}$  w.r.t. the above gradation by  $\mathcal{S}$ .

The natural gradation of the exterior algebra  $\wedge^*(V)$  yields another gradation on  $\mathcal{S}$  as follows:

$$\mathcal{S} = \bigoplus_{p \geq 0} \mathcal{S}^p, \mathcal{S}^p = \mathcal{A} \otimes \wedge^p(V).$$

In other words, elements of  $\mathcal{S}^p$  have the form

$$P = P_{s_1 \dots s_p}^{i_1 \dots i_p} \theta_{i_1}^{s_1} \dots \theta_{i_p}^{s_p} + X_{t_1 \dots t_{p-1}}^{j_1 \dots j_{p-1}} \zeta \theta_{j_1}^{t_1} \dots \theta_{j_{p-1}}^{t_{p-1}},$$

where  $P_{s_1 \dots s_p}^{i_1 \dots i_p}, X_{t_1 \dots t_{p-1}}^{j_1 \dots j_{p-1}} \in \mathcal{A}$ , these coefficients are uniquely determined by  $P$  if we require that they are antisymmetric w.r.t. the exchange of the indices  $(i_k, s_k) \leftrightarrow (i_l, s_l)$  and  $(j_k, t_k) \leftrightarrow (j_l, t_l)$ . An element  $P \in \mathcal{S}$  is called local if it does not depend on  $\zeta$ . The subspace of  $\mathcal{S}$  consists of local elements is denoted by  $\mathcal{S}_{loc}$

**Definition 2.3.2** We introduce a derivation  $\hat{\partial} : \mathcal{S} \rightarrow \mathcal{S}$

$$\hat{\partial} = u^{i, s+1} \partial_{i, s} + \theta_i^{s+1} \partial_s^i - u^{i, 1} \theta_i \partial_\zeta, \quad (2.10)$$

and denote by  $\mathcal{E}$  the quotient space  $\mathcal{S}/\hat{\partial}\mathcal{S}$ . Here  $\partial_{i, s}, \partial_s^i, \partial_\zeta$  are derivations on  $\mathcal{S}$  given by

$$\partial_{i, s} = \frac{\partial}{\partial u^{i, s}}, \quad \partial_s^i = \frac{\partial}{\partial \theta_i^s}, \quad \partial_\zeta = \frac{\partial}{\partial \zeta}.$$

The coset of  $P \in \mathcal{S}$  in  $\mathcal{E}$  is denoted by  $\int_{\mathbb{R}} P dx$ , and  $P$  is called a density of  $\int_{\mathbb{R}} P dx$ . A coset  $\int_{\mathbb{R}} P dx \in \mathcal{E}$  is called local if it possesses a local density  $P \in \mathcal{S}_{loc}$ . We denote the subspace of  $\mathcal{E}$  consists of local elements by  $\mathcal{E}_{loc}$ . Since  $\hat{\partial}$  is homogeneous w.r.t. both gradations of  $\mathcal{S}$ , the space  $\mathcal{E}$  also has two gradations.

Note that the restriction of  $\hat{\partial}$  on  $\mathcal{A} = \mathcal{S}^0$  coincides with  $\partial$ , so it is not necessary to distinguish  $\partial$  and  $\hat{\partial}$ . We will denote  $\hat{\partial}$  by  $\partial$  henceforth.

The main result of the present subsection is the following theorem.

**Theorem 2.3.3** The space  $\mathcal{E}$  is isomorphic to the space of quasi-local multi-vectors.

*Proof* We first define a map  $j : \mathcal{S}^p \rightarrow \text{Alt}^p(\mathcal{A}, \mathcal{A})$  as follow:

$$\begin{aligned} j(P)(f_1, \dots, f_p) &= \partial_{s_p}^{i_p} \dots \partial_{s_1}^{i_1} (P) \cdot \partial^{s_1} \delta_{i_1}(f_1) \dots \partial^{s_p} \delta_{i_p}(f_p) \\ &+ \partial_{t_{p-1}}^{j_{p-1}} \dots \partial_{t_1}^{j_1} \partial_\zeta(P) \cdot \prod_{k=1}^p (-1)^{k-1} E(f_k) \partial^{t_1} \delta_{j_1}(f_1) \dots \hat{f}_k \dots \partial^{t_{p-1}} \delta_{j_{p-1}}(f_{p-1}). \end{aligned} \quad (2.11)$$

The identities  $\partial_s^i \partial_t^j + \partial_t^j \partial_s^i = 0$ ,  $\partial_s^i \partial_\zeta + \partial_\zeta \partial_s^i = 0$  imply that  $j(P)$  is alternating. Since  $\delta_i \partial = 0$ ,  $E \partial = 0$ , the map  $j$  induces a map  $j' : \mathcal{S}^p \rightarrow \text{Alt}^p(\mathcal{F}, \mathcal{A})$ .

The desired isomorphism is defined as  $j'' : \mathcal{E}^p \rightarrow \mathcal{V}^p$

$$\int_{\mathbb{R}} j'' \quad P dx \quad (F_1, \dots, F_p) = \int_{\mathbb{R}} j'(P)(F_1, \dots, F_p) dx, \quad (2.12)$$

This definition is independent of the choice of the density  $P$ , since we have

$$j(\partial Q)(f_1, \dots, f_p) = \partial(j(Q)(f_1, \dots, f_p)),$$

where  $Q \in \mathcal{S}$  and  $f_1, \dots, f_p \in \mathcal{A}$ .

It is easy to see that the image of  $j''$  is just the subspace of quasi-local multi-vectors, the nontrivial part of the theorem is to prove that  $j''$  is injective, which is equivalent to the following lemma:

**Lemma 2.3.4** *If for  $P \in \mathcal{S}^p$  and any  $f_1, \dots, f_p \in \mathcal{A}$  we have*

$$j(P)(f_1, \dots, f_p) \sim 0,$$

*then there exists  $Q \in \mathcal{S}$  such that  $P = \partial Q$ .*

The proof of the lemma will be given below, the theorem is then proved.  $\square$

In order to prove Lemma 2.3.4, we first need to prove the following two lemmas.

**Lemma 2.3.5** *Lemma 2.3.4 holds true when  $p = 0, 1, 2$ .*

*Proof* The case with  $p = 0$  is trivial. We begin with  $p = 1$ . Let  $P \in \mathcal{S}^1$ , so

$$P = Y_s^i \theta_i^s + h \zeta \sim X^i \theta_i + h \zeta,$$

where  $X^i = \sum_{s \geq 0} (-\partial)^s Y_s^i$ . Let  $f \in \mathcal{A}$ , then

$$j(P)(f) \sim X^i \delta_i(f) + h E(f) \sim 0.$$

Take  $f = 1$ , we obtain  $h \sim 0$ , so there exists  $h' \in \mathcal{A}$  such that  $h = \partial h'$ , then

$$P \sim X^i \theta_i + h \zeta \sim Z^i \theta_i,$$

where  $Z^i = X^i + h' u^{i,1}$ , and

$$j(P)(f) \sim Z^i \delta_i(f) \sim 0.$$

According to Lemma 2.1.7,  $Z^i = c u^{i,1}$ , so

$$P \sim c u^{i,1} \theta_i = \partial(-c \zeta) \sim 0.$$

The  $p = 1$  case is proved.

Now let  $P \in \mathcal{S}^2$ , by performing integration by parts we have

$$P \sim P_s^{ij} \theta_i^s \theta_j + X^i \zeta \theta_i,$$

where  $P_s^{ij}$  are uniquely determined by the antisymmetric condition

$$P_s^{ij} + \sum_{t \geq s} (-1)^t \binom{t}{s} \partial^{t-s} P_t^{ji} = 0.$$

Then

$$j(P)(f, g) \sim 2P_s^{ij} \partial^s \delta_i(f) \delta_j(g) + X^i (E(f) \delta_i(g) - \delta_i(f) E(g)) \sim 0.$$

When  $f = 1$  we have  $X^i \delta_i(g) \sim 0$ , so  $X^i = c u^{i,1}$ , thus

$$j(P)(f, g) \sim 2P_s^{ij} \partial^s \delta_i(f) + 2c u^{j,1} E(f) \delta_j(g) \sim 0,$$

so there exists a differential operator  $\tilde{c} : \mathcal{A} \rightarrow \mathbb{R}$  such that

$$2P_s^{ij} \partial^s \delta_i(f) + 2c u^{j,1} E(f) = \tilde{c}(f) u^{j,1},$$

but Lemma 2.1.4 tells us such  $\tilde{c}$  must vanish, so

$$2P_s^{ij} \partial^s \delta_i(f) + 2c u^{j,1} E(f) = 0,$$

then choose  $f$  to be the monomials  $1, u^i, u^i u^j, \dots$ , one can obtain  $P_s^{ij} = 0$  and  $c = 0$ , so  $P \sim 0$ . The lemma is proved.  $\square$

**Lemma 2.3.6** *Let  $p \geq 2$ ,  $P \in \mathcal{S}_{loc}^p$ , if for any  $f_1, \dots, f_p \in \mathcal{A}$ ,*

$$j(P)(f_1, \dots, f_p) \sim 0,$$

*then there exists  $Q \in \mathcal{S}_{loc}^p$  such that  $P = \partial Q$ .*

*Proof* Note that  $P$  is homogeneous function of  $\theta_i^s$ , so

$$P = \frac{1}{p} \theta_i^s \partial_s^i(P) \sim \frac{1}{p} \theta_i \delta^i(P) = \frac{1}{p} N(P), \quad (2.13)$$

where

$$\delta^i = \prod_{s \geq 0} (-\partial)^s \partial_s^i,$$

and

$$N = \theta_i \delta^i \quad (2.14)$$

is the normalizing operator used in [5].

By the definitions of  $j$  and  $\delta^i$ , it is easy to see that

$$j(P)(f_1, \dots, f_p) \sim j(\delta^i(P))(f_2, \dots, f_p) \delta_i(f_1) \sim 0,$$

so we have

$$j(\delta^i(P))(f_2, \dots, f_p) = c(f_2, \dots, f_p) u^{i,1},$$

where  $c(f_2, \dots, f_p) \in \mathbb{R}$  for any  $f_2, \dots, f_p$ . According to Lemma 2.1.4, the map  $j(\delta^i(P))$  is zero (here we used the condition  $p \geq 2$ ). By choosing suitable  $f_2, \dots, f_p$ , one can obtain  $\delta^i(P) = 0$ , so from (2.13) it follows that  $P \sim 0$ . The lemma is proved.  $\square$

**Remark 2.3.7** *The above lemma is not correct when  $p = 1$ . In fact,  $P = u^{i,1} \theta_i$  satisfies the condition, but  $P = -\partial \zeta$ , and  $\zeta \notin \mathcal{S}_{loc}^1$ . This fact also explains why the super variable  $\zeta$  is so important.*

*Proof of Lemma 2.3.4* Since the cases with  $p = 0, 1, 2$  have been proved, we assume  $p \geq 3$ . It is easy to see

$$j(P)(1, f_2, \dots, f_p) = -j(\partial_\zeta(P))(f_2, \dots, f_p) \sim 0,$$

the derivative  $\partial_\zeta(P)$  satisfies the condition of Lemma 2.3.6, so we have

$$\partial_\zeta(P) = \partial Q, \text{ where } Q \in \mathcal{S}_{loc}^p.$$

Let  $P_0 = P - \zeta \partial_\zeta(P)$ , then

$$P = P_0 + \zeta \partial_\zeta(P) = P_0 + \zeta \partial(Q) \sim P_0 + u^{i,1} \theta_i Q \in \mathcal{S}_{loc}^p.$$

By applying Lemma 2.3.6 again, we proved the lemma.  $\square$

Due to Theorem 2.3.3, we denote the space of quasi-local multi-vectors also by  $\mathcal{E}$ .

## 2.4 The Schouten-Nijenhuis bracket

In this subsection we define the Schouten-Nijenhuis bracket on the space  $\mathcal{E}$  of quasi-local multi-vectors, it turns out to be the restriction of the Nijenhuis-Richardson bracket defined on the space  $\mathcal{V}$  of generalized multi-vectors to  $\mathcal{E}$ .

The original Nijenhuis-Richardson bracket was defined in [46, 47] for finite dimensional vector spaces. In [6], this construction is generalized to vector spaces that are not necessarily finite dimensional. In what follows, we recall the definition of this construction, and present it in an axiomatic way.

**Theorem 2.4.1** *There exists a unique bilinear map  $[\cdot, \cdot] : \mathcal{V}^p \times \mathcal{V}^q \rightarrow \mathcal{V}^{p+q-1}$  satisfying the following conditions:*

$$[P, F](F_2, \dots, F_p) = P(F, F_2, \dots, F_p), \quad (2.15)$$

$$[P, Q] = (-1)^{pq}[Q, P], \quad (2.16)$$

$$[[P, Q], F] + (-1)^{qp}[[Q, F], P] + [[F, P], Q] = 0 \quad (2.17)$$

for any  $P \in \mathcal{V}^p$ ,  $Q \in \mathcal{V}^q$ ,  $F, F_2, \dots, F_p \in \mathcal{F}$ . It is called the Nijenhuis-Richardson bracket of the generalized multi-vectors over  $\mathcal{F}$ .

*Proof* Note that for  $P \in \mathcal{V}^p$ ,  $Q \in \mathcal{V}^q$ ,  $F_1, F_2, \dots, F_{p+q-1} \in \mathcal{F}$ ,

$$\begin{aligned} [P, Q](F_1, F_2, \dots, F_{p+q-1}) &= [[P, Q], F_1](F_2, \dots, F_{p+q-1}) \\ &= -((-1)^{qp}[[Q, F_1], P] + [[F_1, P], Q])(F_2, \dots, F_{p+q-1}), \end{aligned}$$

so the bracket on  $\mathcal{V}^p \times \mathcal{V}^q$  is determined by the brackets on  $\mathcal{V}^{p-1} \times \mathcal{V}^q$  and  $\mathcal{V}^p \times \mathcal{V}^{q-1}$ , thus the uniqueness can be proved by induction on  $p + q$  immediately.

To prove the existence, we recall the product  $\bar{\wedge} : \mathcal{V}^p \times \mathcal{V}^q \rightarrow \mathcal{V}^{p+q-1}$  defined in [47]:

$$P \bar{\wedge} Q(F_1, \dots, F_{p+q-1}) = \sum_{I \in S_{p,q}} (-1)^{|I|} P(Q(F_{i_1}, \dots, F_{i_q}), F_{i_{q+1}}, \dots, F_{i_{p+q-1}}),$$

where  $S_{p,q}$  is the subset of the symmetry group  $S_{p+q-1}$

$$S_{p,q} = \{ I = (i_1, \dots, i_{p+q-1}) \in S_{p+q-1} \mid \begin{array}{l} i_1 < i_2 < \dots < i_q \\ i_{q+1} < i_{q+2} < \dots < i_{p+q-1} \end{array} \},$$

and  $|I|$  denotes the parity of the permutation  $I$ . Note that this product is neither commutative nor associative.

We define the desired bracket as

$$[P, Q] = (-1)^{(p+1)q} P \bar{\wedge} Q + (-1)^p Q \bar{\wedge} P. \quad (2.18)$$

It is easy to verify that (2.18) satisfies the conditions (2.15) and (2.16). So we only need to prove the validity of the condition (2.17).

Let  $P \in \mathcal{V}^p$ ,  $Q \in \mathcal{V}^q$ ,  $F_1 \in \mathcal{F}$ . Denote

$$\tilde{P} = [P, F_1], \quad \tilde{Q} = [Q, F_1].$$

By definition of the product  $\bar{\wedge}$ , we have

$$\begin{aligned} & P \bar{\wedge} Q(F_1, F_2, \dots, F_{p+q-1}) \\ &= P \bar{\wedge} \tilde{Q}(F_2, \dots, F_{p+q-1}) + (-1)^{q+1} \tilde{P} \bar{\wedge} Q(F_2, \dots, F_{p+q-1}). \end{aligned} \quad (2.19)$$

It yields the identity

$$[[P, Q], F_1] = -(-1)^p [P, \tilde{Q}] - [\tilde{P}, Q],$$

which is equivalent to (2.17). The theorem is proved.  $\square$

**Corollary 2.4.2** *The Nijenhuis-Richardson bracket satisfies the following graded Jacobi identity*

$$(-1)^{pr} [[P, Q], R] + (-1)^{qp} [[Q, R], P] + (-1)^{rq} [[R, P], Q] = 0, \quad (2.20)$$

for any  $P \in \mathcal{V}^p$ ,  $Q \in \mathcal{V}^q$ ,  $R \in \mathcal{V}^r$ .

*Proof* We prove the corollary by induction on  $p + q + r$ . When  $r = 0$ , it is just the condition (2.17), when  $r > 0$ , we assume that the condition (2.20) holds true for any  $p', q', r'$  with  $p' + q' + r' < p + q + r$ . Let  $P \in \mathcal{V}^p$ ,  $Q \in \mathcal{V}^q$ ,  $R \in \mathcal{V}^r$ ,  $F \in \mathcal{F}$ , then

$$\begin{aligned} & (-1)^{pr} [[[P, Q], R], F] \\ &= (-1)^{pr+p+q} [[P, Q], \tilde{R}] + (-1)^{pr+p} [[P, \tilde{Q}], R] + (-1)^{pr} [[\tilde{P}, Q], R], \end{aligned}$$

where  $\tilde{K} = [K, F]$ ,  $K = P, Q, R$ , so we have

$$[(-1)^{pr} [[P, Q], R] + (-1)^{qp} [[Q, R], P] + (-1)^{rq} [[R, P], Q], F] = 0.$$

Thus the corollary follows from the property (2.15).  $\square$

It is a nontrivial fact that the space  $\mathcal{E}$  of quasi-local multi-vectors is closed under the operation of Richardson-Nijenhuis bracket. In what follows we first define a bracket on  $\mathcal{E}$  by using the super variable description of the space of quasi-local multi-vectors, and we prove that this bracket satisfies the conditions of Theorem 2.4.1. Then the uniqueness property of Theorem 2.4.1 shows that this bracket coincides with the restriction of the Nijenhuis-Richardson bracket to  $\mathcal{E}$ . To this end, let us first introduce some notations.

**Definition 2.4.3** For  $i = 1, \dots, n$ , and integers  $\alpha, s \geq 0$ , we define the higher generalized momentum operator, the higher energy operator and the energy operator from  $\mathcal{S}$  to  $\mathcal{S}$  as

$$\begin{aligned} p_{i,\alpha,s} &= \sum_{\beta \geq 0} (-1)^\beta \binom{\beta+s}{s} \partial^\beta \partial_{i,\alpha+\beta+s}, \\ p_{\alpha,s}^i &= \sum_{\beta \geq 0} (-1)^\beta \binom{\beta+s}{s} \partial^\beta \partial_{\alpha+\beta+s}^i, \\ E_s &= \sum_{\alpha \geq 1} u^{i,\alpha} p_{i,\alpha,s} + \theta_i^\alpha p_{\alpha,s}^i, \\ E &= E_0 - 1, \end{aligned}$$

and assume that  $p_{i,\alpha,-1} = \partial_{i,\alpha-1}$ ,  $p_{\alpha,-1}^i = \partial_{\alpha-1}^i$ . In particular, we have

$$E_{-1} = \sum_{\alpha \geq 1} u^{i,\alpha} \partial_{i,\alpha-1} + \theta_i^\alpha \partial_{\alpha-1}^i = \partial + u^{i,1} \theta_i \partial_\zeta,$$

and the higher Euler operators from  $\mathcal{S}$  to  $\mathcal{S}$  is defined as

$$\delta_{i,s} = p_{i,0,s}, \quad \delta_s^i = p_{0,s}^i, \quad \delta_i = \delta_{i,0}, \quad \delta^i = \delta_0^i. \quad (2.21)$$

There is another useful operator  $\hat{E} : \mathcal{S} \rightarrow \mathcal{S}$ , which is defined as

$$\hat{E} = E + N, \quad (2.22)$$

where  $N$  is the normalizing operator defined in (2.14).

**Theorem 2.4.4** Define the bilinear map  $[\cdot, \cdot] : \mathcal{E}^p \times \mathcal{E}^q \rightarrow \mathcal{E}^{p+q-1}$  by

$$\begin{aligned} & \int \int [P dx, Q dx] \\ &= \int \delta^i(P) \delta_i(Q) + (-1)^p \delta_i(P) \delta^i(Q) + \partial_\zeta(P) \hat{E}(Q) + (-1)^p \hat{E}(P) \partial_\zeta(Q) dx. \end{aligned} \quad (2.23)$$

It satisfies the condition (2.15)-(2.17), so it coincides with the Nijenhuis-Richardson bracket defined on  $\mathcal{V}$ .

We call the above bracket (2.23) the *Schouten-Nijenhuis bracket* among quasi-local multi-vectors.

**Corollary 2.4.5** *The bracket (2.23) satisfies the graded Jacobi identity*

$$(-1)^{pr}[[P, Q], R] + (-1)^{qp}[[Q, R], P] + (-1)^{rq}[[R, P], Q] = 0, \quad (2.24)$$

for any  $P \in \mathcal{E}^p$ ,  $Q \in \mathcal{E}^q$ ,  $R \in \mathcal{E}^r$ .

**Remark 2.4.6** *The Schouten-Nijenhuis bracket can be restricted on the subspace  $\mathcal{E}_{loc}$ , the resulting bracket among local multi-vectors is equivalent to the brackets defined by Getzler [29] and by Kersten, Krasil'shchik, Verbovetsky [32].*

**Remark 2.4.7** *If  $P, Q$  are of degree zero, then they become pairs of multi-vectors on the manifold  $M$ , and the bracket (2.23) degenerates to*

$$\begin{aligned} [P, Q] &= \partial^i(P)\partial_i(Q) + (-1)^p\partial_i(P)\partial^i(Q) \\ &\quad + (1-q)\partial_\zeta(P)Q + (1-p)(-1)^pP\partial_\zeta(Q). \end{aligned}$$

*This formula has appeared in the study of classical Jacobi structures, see [31] for example.*

*Proof of Theorem 2.4.4* We first show that the bracket is well-defined, i.e. it is independent of the choices of  $P, Q \in \mathcal{S}$ . Introduce a bilinear map  $[\ , ]_{pr} : \mathcal{S}^p \times \mathcal{S}^q \rightarrow \mathcal{S}^{p+q-1}$

$$[P, Q]_{pr} = \delta^i(P)\delta_i(Q) + (-1)^p\delta_i(P)\delta^i(Q) + \partial_\zeta(P)\hat{E}(Q) + (-1)^p\hat{E}(P)\partial_\zeta(Q),$$

then it is easy to see that  $[P, Q]_{pr} \sim D_P(Q)$ , where  $D_P : \mathcal{S} \rightarrow \mathcal{S}$  is a first order differential operator

$$D_P = -\partial_\zeta(P) + (-1)^p\hat{E}(P)\partial_\zeta + X^{i,s}\partial_{i,s} + Y_i^s\partial_s^i, \quad (2.25)$$

$X^{i,s}, Y_i^s$  are defined by

$$X^{i,s} = \partial^s X^i + \bigotimes_{t=1}^s \partial^{s-t} \partial_\zeta(P)u^{i,t}, \quad Y_i^s = \partial^s(Y_i) + \bigotimes_{t=1}^s \partial^{s-t} \partial_\zeta(P)\theta_i^t,$$

and  $X^i = \delta^i(P)$ ,  $Y_i = (-1)^p(\delta_i(P) - \theta_i\partial_\zeta(P))$ .

Note that the operator  $D_P$  satisfies  $D_P\partial = \partial(D_P - D_P(1))$ , so

$$[P, \partial(Q)]_{pr} \sim D_P\partial(Q) = \partial(D_P - D_P(1))(Q) \sim 0,$$

which implies that the definition is independent of the choice of  $Q$ . Then by using the fact that

$$[P, Q]_{pr} = (-1)^{pq}[Q, P]_{pr},$$

it is easy to see that this definition is also independent of the choice of  $P$ , so it is well-defined on  $\mathcal{E}^p \times \mathcal{E}^q$ .

We next verify that the bracket defined in (2.23) satisfies the conditions (2.15)-(2.17). In fact, the conditions (2.15) and (2.16) are easy to verify, so we omit their proof here. The condition (2.17) is a consequence of the following lemma:

**Lemma 2.4.8** *Define*

$$Z_{P,f} = D_{D_f(P)} + D_f D_P + (-1)^p D_P D_f \quad (2.26)$$

for  $P \in \mathcal{S}^p$ ,  $f \in \mathcal{A}$ , then  $Z_{P,f} = 0$ .

The theorem then follows from this lemma, which will be proved below.  $\square$

In order to prove Lemma 2.4.8, we first need to prove some other lemmas.

**Lemma 2.4.9** *For  $X, Y \in \mathcal{E}^1$ ,  $F \in \mathcal{F}$ , we have*

$$[X, Y](F) = X(Y(F)) - Y(X(F)),$$

here  $X(\cdot)$  means  $j''(X)(\cdot)$ , we omit  $j''$  from now on.

*Proof* The general form of an element in  $\mathcal{E}^1$  reads

$$X = \int X^i \theta_i + a \zeta \quad dx,$$

so we need to prove the lemma for the following three cases:

1.  $X = \int (X^i \theta_i) \quad dx$ ,  $Y = \int (Y^j \theta_j) \quad dx$ ;
2.  $X = \int (X^i \theta_i) \quad dx$ ,  $Y = \int (b \zeta) \quad dx$ ;
3.  $X = \int (a \zeta) \quad dx$ ,  $Y = \int (b \zeta) \quad dx$ ;

We give below the proof for the third case, the proofs for the other two cases are similar.

According to the definition of  $[ \quad ]$ , we have

$$\begin{aligned} [X, Y] &= \int a \hat{E}(b \zeta) - b \hat{E}(a \zeta) \quad dx \\ &= \int \int (-1)^s a u^{i,t} \partial^s \partial_{i,s+t} (b \zeta) - b u^{i,t} \partial^s \partial_{i,s+t} (a \zeta) \quad dx \\ &= \int \int_{s \geq 0} \int_{t \geq 1} \partial^s a u^{i,t} b_{(i,s+t)} - \partial^s b u^{i,t} a_{(i,s+t)} \quad \zeta \quad dx \\ &= \int_{t \geq 0} \int_{s \geq 0} \int_{t \geq 1} \partial^t (a E_t(b)) - \partial^t (b E_t(a)) \quad \zeta \quad dx. \end{aligned}$$

Here the last equality follows from the last identity of Lemma 2.1.3. In what follows we will also use this lemma. So the left hand side reads (we omit the integral symbol to save space below)

$$[X, Y](F) \sim \int_{t \geq 0} \partial^t (a E_t(b)) - \partial^t (b E_t(a)) \quad E(F).$$

On the other hand, let  $F = \int_{\mathbb{R}} f dx$ , we have

$$\begin{aligned}
& (XY - YX)(F) \sim a E(b E(f)) - b E(a E(f)) \\
= & \sum_{t \geq 0} a(-1)^t E_t(b) \partial^t E(f) + \partial^t(b) E_t E(f) - a b E(f) \\
& - \sum_{t \geq 0} b(-1)^t E_t(a) \partial^t E(f) + \partial^t(a) E_t E(f) + a b E(f) \\
\sim & \sum_{t \geq 0} \partial^t (a E_t(b)) - \partial^t (b E_t(a)) E(F) \\
& + b \sum_{t \geq 0} \partial^t (a E_t E(f)) - \sum_{t \geq 0} (-1)^t \partial^t(a) E_t E(f) \quad !,
\end{aligned}$$

so we only need to prove the following identity

$$\sum_{t \geq 0} \partial^t (a E_t E(f)) = \sum_{t \geq 0} (-1)^t \partial^t(a) E_t E(f),$$

which is an easy corollary of the identity iii) in Lemma 2.1.5. The lemma is proved.  $\square$

The following lemma is a generalization of Lemma 2.1.9.

**Lemma 2.4.10** *Let  $f \in \mathcal{S}$ , if for any  $g \in \mathcal{A}$  we have  $fg \sim 0$ , then  $f = 0$ .*

*Proof* We first prove the lemma when  $f$  is local. Without loss of generality, we assume that  $f \in \mathcal{S}_{loc}^p$  and is homogeneous w.r.t. the gradation of  $\mathcal{S}^p$  defined in (2.9). The condition of the lemma can be written as

$$gf = \partial(h). \tag{2.27}$$

The  $p = 0$  case is just Lemma 2.1.9. Now we assume  $p = 1$ , so that

$$f = \sum_{s=0}^{\infty} f_s^i \theta_i^s, \quad h = \sum_{s=0}^{\infty} h_s^i \theta_i^s + a \zeta.$$

Here  $h$  is also homogeneous w.r.t. to both gradation of  $\mathcal{S}$ . Compare the coefficients of  $\theta_i^s$  and  $\zeta$  on both sides of (2.27) we obtain

$$\partial(a) = 0, \quad g f_0^i = \partial h_0^i - a u^{i,1}, \quad g f_s^i = h_{s-1}^i + \partial h_s^i \quad (s \geq 1), \tag{2.28}$$

from which we obtain  $a \in \mathbb{R}$  and  $g f_0^i \sim 0$ . By applying Lemma 2.1.9, we also see that  $f_0^i = 0$  and  $h_0^i = a u^i$ . From (2.28) it also follows that

$$a = a(g) = \frac{1}{u^i} \sum_{l=0}^{\infty} (-\partial)^{l-1} f_{l+1}^i \quad !,$$

so by using Lemma 2.1.4 we see that  $a = 0$  and thus  $h_0^i = 0$ . Now by using the third equation of (2.28), Lemma 2.1.9 and the homogeneity of  $h$  we arrive at  $f = 0$ . Thus we proved the case when  $p = 1$ .

When  $p \geq 2$ , let  $f g = \partial(h)$  for certain  $h \in \mathcal{S}^2$ . We claim that  $h \in \mathcal{S}_{loc}^p$ . In fact if  $h = h_0 + \zeta h_1$  with  $h_0, h_1 \in \mathcal{S}_{loc}$ , then  $\partial(h) = f g \in \mathcal{S}_{loc}$ , so  $\partial(h_1) = 0$  which implies that  $h_1$  is a constant, and thus  $\zeta h_1 \in \mathcal{S}^1$ . Since  $h \in \mathcal{S}^2$  we must have  $h_1 = 0$ . Now for the local multi-vector  $h \in \mathcal{S}_{loc}^p$  one can prove that  $\delta^i \partial(h) = 0$ , so

$$0 = \delta^i (f g) = \sum_{t \geq 0} (-1)^t \delta_t^i(f) \partial^t(g). \quad (2.29)$$

Since we have assumed that  $f$  is homogeneous in both gradations of  $\mathcal{S}$ , there are only finite number of nonzero terms in the right hand side of the above equation. We regard the equation (2.29) as a system of linear equations of  $(-1)^t \delta_t^i(f)$ , then by choosing linear independent  $g$ 's and using the property of Wronskian determinants, we obtain  $\delta_t^i(f) = 0$  for all  $t \geq 0$ . Then the following identities

$$f = \frac{1}{p} \theta_i^s \partial_s^i(f), \quad \partial_t^i(f) = \sum_{s \geq 0} \binom{s+t}{s} \partial^s \delta_{s+t}^i(f)$$

imply that  $f = 0$ . Thus we proved the case when  $f$  is local.

For the nonlocal case we note that  $[\partial, \partial_\zeta] = 0$ , so

$$f g \sim 0 \quad \Rightarrow \quad \partial_\zeta(f) g \sim 0,$$

by applying the above result to the local  $\partial_\zeta f$  we obtain  $\partial_\zeta(f) = 0$ , so  $f \in \mathcal{S}_{loc}$ . By applying the above result again to  $f$  we finish the proof of the lemma.  $\square$

**Remark 2.4.11** *The identity  $\delta^i \partial(h) = 0$  is not true if  $\partial_\zeta h \neq 0$ . In general, the identities listed in Lemma 2.1.3 should be modified when the operators act on  $\mathcal{S}$ .*

*Proof of Lemma 2.4.8* It is easy to see that the operator

The last expression equals zero since by using the fact that  $[D_f, \partial] = 0$  we obtain

$$\delta^i D_f + D_f \delta^i = \sum_{s \geq 0} (-\partial)^s \partial_s^i D_f + D_f \partial_s^i = 0.$$

For the other two identities, we first consider the case when  $p = 1$ . Let  $P \in \mathcal{S}^1$  and  $Y = Y^i \theta_i + h \zeta \in \mathcal{S}^1$ , Lemma 2.4.9 implies

$$0 = \int Z_{P,f}(Y) dx = \int Z_{P,f}(\zeta)h + Z_{P,f}(\theta_i)Y^i dx,$$

then Lemma 2.1.9 implies that  $Z_{P,f}(\zeta) = 0$  and  $Z_{P,f}(\theta_i) = 0$ , so we have  $Z_{P,f} = 0$ .

Finally for  $P \in \mathcal{S}^p$  and  $Y = Y^i \theta_i + h \zeta \in \mathcal{S}^1$  we have

$$0 = \int (-1)^p Z_{Y,f}(P) dx = \int Z_{P,f}(Y) dx.$$

By using the same argument as we used for the  $p = 1$  case and by using Lemma 2.4.10 we obtain  $Z_{P,f}(\zeta) = 0$  and  $Z_{P,f}(\theta_i) = 0$ , so we have  $Z_{P,f} = 0$ .

The lemma is proved. □

## 2.5 Miura type transformations

On the differential algebra  $(\mathcal{A}, \partial)$  there is a natural class of coordinate transformations which are called Miura type transformations in [21], they induce transformations of the graded Lie algebra  $(\mathcal{E}, [ , ])$ . We are to give some useful formulae for these transformations in this subsection.

**Definition 2.5.1** *Let  $\bar{u}^1, \dots, \bar{u}^n \in \mathcal{A}$  be  $n$  differential polynomials, and  $\bar{u}_0^i$  be the degree zero components of  $\bar{u}^i$ , if*

$$\det \frac{\partial \bar{u}_0^i}{\partial u^j} \neq 0$$

*then the map  $(u^1, \dots, u^n) \mapsto (\bar{u}^1, \dots, \bar{u}^n)$  defines a coordinate transformation on  $\mathcal{A}$  which is called a Miura type transformation. When  $\bar{u}^i = \bar{u}_0^i$ , this transformation is called a Miura type transformation of the first kind; when  $\bar{u}_0^i = u^i$ , this transformation is called a Miura type transformation of the second kind.*

It is easy to see that every Miura type transformation is the composition of a first kind Miura type transformation and a second kind one. A first kind Miura type transformation is just a change of local coordinates on the base manifold  $M$ , and a general Miura type transformation is an analogue of change of local coordinates on the jet space  $J^\infty(M)$ , note that  $\mathcal{A}$  is similar to but different from  $C^\infty(J^\infty(M))$ .

Given a Miura type transformation  $(\bar{u}^i)$ , we can define  $\bar{u}^{i,s} = \partial^s (\bar{u}^i)$ , and express  $u^i$  as differential polynomials in  $(\bar{u}^{i,s})$ , then every element of  $\mathcal{A}$  can be expressed as differential polynomial in  $(\bar{u}^{i,s})$ , so we can define the following derivation

$$\bar{\partial}_{i,s} = \frac{\partial}{\partial \bar{u}^{i,s}} : \mathcal{A} \rightarrow \mathcal{A},$$

and the derivation  $\partial : \mathcal{A} \rightarrow \mathcal{A}$  now reads

$$\partial = \sum_{s \geq 0} \bar{u}^{i,s+1} \bar{\partial}_{i,s}.$$

Furthermore, one can define operators  $\bar{p}_{i,\alpha,s}$ ,  $\bar{E}_s$  and  $\bar{E}$  by replacing  $\partial_{i,s}$ ,  $u^{i,s}$  by  $\bar{\partial}_{i,s}$ ,  $\bar{u}^{i,s}$  respectively in the original definition of  $p_{i,\alpha,s}$ ,  $E_s$ ,  $E$  given in Section 2.

**Lemma 2.5.2** *Let  $f \in \mathcal{A}$  be a differential polynomial, then the following identities hold true:*

$$\delta_i(f) = \sum_{s \geq 0} (-\partial)^s \partial_{i,s}(\bar{u}^j) \bar{\delta}_j(f), \quad (2.30)$$

$$E(f) = \bar{E}(f) - \sum_{s \geq 0} \sum_{\alpha \geq 1} u^{i,\alpha} (-\partial)^s \partial_{i,s+\alpha}(\bar{u}^j) \bar{\delta}_j(f). \quad (2.31)$$

*Proof* We denote the right hand sides of (2.30) and (2.31) by  $g_i$  and  $h$  respectively. According to Corollary 2.1.10, we only need to prove that for any first order differential operator  $D : \mathcal{A} \rightarrow \mathcal{A}$  satisfying  $D\partial = \partial(D - D(1))$ , one has

$$D(f) \sim X^i g_i - a h, \quad (2.32)$$

where  $a = D(1)$  and  $X^i = (D - D(1))(u^i)$ .

On the other hand, if we apply Lemma 2.1.8 in the coordinate system  $(\bar{u}^{i,s})$ , we obtain

$$D(f) \sim \bar{X}^j \bar{\delta}_j(f) - \bar{a} \bar{E}(f), \quad (2.33)$$

where  $\bar{a} = D(1) = a$ , and

$$\bar{X}^j = (D - D(1))(\bar{u}^j) = \sum_{s \geq 0} \partial^s (X^i) - \sum_{t=1}^{\infty} \partial^{s-t} (a u^{i,t}) \bar{\partial}_{i,s}(\bar{u}^j).$$

Then one can obtain (2.32) from (2.33) via integration by parts.  $\square$

**Example 2.5.3** *Let  $(\bar{u}^i)$  be a Miura type transformation of the first kind, then the identities (2.30) and (2.31) read*

$$\delta_i(f) = \partial_{i,0}(\bar{u}^j) \bar{\delta}_j(f), \quad E(f) = \bar{E}(f).$$

*The first one gives the classical transformation formula of variational derivatives, while the second one means that the energy is independent of the choice of coordinates system.*

If we regard  $\delta_i$  and  $E$  as super variables  $\theta_i$  and  $\zeta$ , the above lemma gives the transformation rule of super variables under Miura type transformations. More precisely, we introduce a new family of super variables  $\bar{\theta}_i^s, \bar{\zeta}$ , and define the space

$$\bar{\mathcal{S}} = \mathcal{A} \otimes \wedge^*(\bar{V}), \text{ where } \bar{V} = \bigoplus_{i,s} \mathbb{R}\bar{\theta}_i^s \oplus \mathbb{R}\bar{\zeta}.$$

We can also define the quotient space  $\bar{\mathcal{E}} = \bar{\mathcal{S}}/\partial\bar{\mathcal{S}}$ , and prove that  $\bar{\mathcal{E}}$  is isomorphic to the space of quasi-local multi-vectors in  $\mathcal{V}$ , so  $\bar{\mathcal{E}}$  is also isomorphic to  $\mathcal{E}$ .

According to Lemma 2.5.2, we introduce the following isomorphism  $\bar{\Phi} : \mathcal{S} \rightarrow \bar{\mathcal{S}}$  of super commutative algebras

$$\begin{aligned} \bar{\Phi}(f) &= f, \quad f \in \mathcal{A}, \\ \bar{\Phi}(\theta_i^s) &= \partial^s \prod_{t \geq 0} (-\partial)^t \partial_{i,t}(\bar{u}^j) \bar{\theta}_j, \\ \bar{\Phi}(\zeta) &= \bar{\zeta} - \prod_{s \geq 0} \prod_{\alpha \geq 1}^{t \geq 0} u^{i,\alpha} (-\partial)^s \partial_{i,s+\alpha}(\bar{u}^j) \bar{\theta}_j. \end{aligned}$$

**Theorem 2.5.4** *Let  $P \in \mathcal{V}^p$  be a quasi-local  $\mathbb{R}$ -vector, suppose the preimages of  $P$  in  $\mathcal{E}$  and  $\bar{\mathcal{E}}$ , under the isomorphism (2.12), are  $\alpha dx$  and  $\bar{\alpha} dx$  respectively, then we have  $\bar{\alpha} \sim \bar{\Phi}(\alpha)$ .*

*Proof* One can prove the theorem by acting  $P$  on arbitrary  $F_1, \dots, F_p \in \mathcal{F}$  and by using Lemma 2.5.2.  $\square$

For a Miura type transformation of the second kind, the formula given in Theorem 2.5.4 is not ease to use. The remaining part of this subsection is devoted to give another simpler formula for the transformation rule of the second kind Miura type transformations. Let us first introduce the following notation:

**Definition 2.5.5** *Let  $F \in \mathcal{S}$  or  $\mathcal{E}$ , one can decompose  $F$  w.r.t. the gradation (2.9) as follows:*

$$F = F_k + F_{k+1} + F_{k+2} + \dots,$$

where  $F_k \neq 0$  and  $\deg F_l = l$ , then the integer  $k$  is called the order of  $F$ , and is denoted by  $\nu(F)$ .

**Lemma 2.5.6** *Let  $(\bar{u}^i)$  be a second kind Miura type transformation, then there exists an  $X \in \mathcal{E}_{loc}^1$  such that  $\nu(X) > 0$ , and*

$$\bar{u}^i = \exp(D_X)(u^i),$$

where  $D_X$  is the first order differential operator associated to  $X$  (see (2.25) for the definition).

*Proof* Suppose  $\bar{u}^i = u^i + F^i$ , where  $\nu(F^i) \geq k > 0$ , let

$$X_{(k)} = \int F_k^i \theta_i dx,$$

then we have

$$\exp(-D_{X_{(k)}})(\bar{u}^i) = u^i + \tilde{F}^i,$$

where  $\nu(\tilde{F}^i) \geq k + 1 > 0$ . So by induction on  $k$ , we can obtain a series

$$X_{(1)}, X_{(2)}, \dots \in \mathcal{E}_{loc}^1,$$

such that

$$\bar{u}^i = \exp(D_{X_{(1)}}) \exp(D_{X_{(2)}}) \dots (u^i).$$

Finally, by using the Baker-Campbell-Hausdorff formula, and note that

$$\nu(X_{(1)}) < \nu(X_{(2)}) < \dots,$$

one can show that there exists  $X \in \mathcal{E}_{loc}^1$  such that  $\nu(X) > 0$ , and

$$\exp(D_X) = \exp(D_{X_{(1)}}) \exp(D_{X_{(2)}}) \dots$$

The lemma is proved. □

From now on, we will denote a second kind Miura type transformation by  $e^X$  for short, where  $X \in \mathcal{E}_{loc}^1$  and satisfies  $\nu(X) > 0$ . When consider Miura type transformations of the second kind, it is convenient to identify  $\mathcal{E}$  and  $\bar{\mathcal{E}}$  and regard the induced isomorphism as automorphism on  $\mathcal{E}$ .

**Theorem 2.5.7** *Let  $e^X$  be a Miura type transformation of the second kind, then the induced automorphism of  $\mathcal{E}$  is given by*

$$\exp(-\text{ad}_X) : \mathcal{E} \rightarrow \mathcal{E},$$

where  $\text{ad}_X = [X, \cdot] : \mathcal{E} \rightarrow \mathcal{E}$  is the adjoint action of  $X$  on  $\mathcal{E}$ .

*Proof* Since both maps are given by exponential, we only need to compute the infinitesimal part of the transformation. For this we replace  $X$  by  $\epsilon X$ , then the Miura type transformation reads

$$\bar{u}^i = u^i + \epsilon X^i + O(\epsilon^2),$$

where  $X = \int X^i \theta_i dx$ .

Let us consider the variation of the elements of  $\mathcal{S}$  under Miura type transformations. For  $f \in \mathcal{A}$ , we need to replace its independent variables  $u^i$  by  $\bar{u}^i - \epsilon X^i + O(\epsilon^2)$  and expand it into power series in  $\epsilon$ , as the result we have

$$f \mapsto f + \epsilon \Delta(f) + O(\epsilon^2) = f - \epsilon \partial^s(X^i) \partial_{i,s}(f) + O(\epsilon^2).$$

For the super variables  $\theta_i^s$  and  $\zeta$ , according to the definition of  $\bar{\Phi}$ , we have

$$\theta_i^s \mapsto \theta_i^s + \epsilon \Delta(\theta_i^s) + O(\epsilon^2), \quad \zeta \mapsto \zeta + \epsilon \Delta(\zeta) + O(\epsilon^2),$$

where

$$\begin{aligned} \Delta(\theta_i^s) &= \partial^s \prod_{t \geq 0} (-\partial)^t \partial_{i,t}(X^j) \theta_j, \\ \Delta(\zeta) &= - \prod_{s \geq 0} \prod_{\alpha \geq 1} u^{i,\alpha} (-\partial)^s \partial_{i,s+\alpha}(X^j) \theta_j. \end{aligned}$$

Now let  $P = \int \alpha dx \in \mathcal{E}$  be a quasi-local multi-vector, then the isomorphism  $\bar{\Phi}$  becomes  $P \mapsto P + \epsilon \Delta(P) + O(\epsilon^2)$ , where

$$\Delta(P) \sim \Delta(u^{i,s}) \partial_{i,s}(\alpha) + \Delta(\theta_i^s) \partial_s^i(\alpha) + \Delta(\zeta) \partial_\zeta(\alpha),$$

then by using Theorem 2.4.4 and integration by parts, one can obtain

$$\Delta(P) = -[X, P].$$

The theorem is proved. □

## 2.6 Reciprocal transformations

In this subsection, we define the reciprocal transformation and give the transformation formula for quasi-local multi-vectors with the help of super variables.

Let  $\rho \in \mathcal{A}$  be an invertible element, we define a new derivation

$$\tilde{\partial} = \rho^{-1} \partial, \tag{2.34}$$

and the new quotient space  $\tilde{\mathcal{F}} = \mathcal{A} / \tilde{\partial} \mathcal{A}$ . The coset of  $\tilde{f} \in \mathcal{A}$  in  $\tilde{\mathcal{F}}$  is denoted by  $\int \tilde{f} d\tilde{x}$ . It is easy to see that there is an isomorphism

$$\Phi_0 : \mathcal{F} \rightarrow \tilde{\mathcal{F}}, \quad \int f dx \mapsto \int \rho^{-1} f d\tilde{x}.$$

We denote  $\tilde{\mathcal{V}} = \text{Alt}^*(\tilde{\mathcal{F}}, \tilde{\mathcal{F}})$ , then one can extend the isomorphism  $\Phi_0 : \mathcal{F} \rightarrow \tilde{\mathcal{F}}$  to an isomorphism  $\Phi : \mathcal{V} \rightarrow \tilde{\mathcal{V}}$  such that

$$\Phi([P, Q]) = [\Phi(P), \Phi(Q)] \tag{2.35}$$

holds true for any  $P, Q \in \mathcal{V}$ . In fact, for any  $P \in \mathcal{V}^p$  the action of  $\Phi(P)$  on  $\tilde{F}_1, \dots, \tilde{F}_p \in \tilde{\mathcal{F}}$  is given as follows:

$$\Phi(P)(\tilde{F}_1, \dots, \tilde{F}_p) = \Phi_0 \int P \Phi_0^{-1} \tilde{F}_1, \dots, \Phi_0^{-1} \tilde{F}_p.$$

By the definition of  $\bar{\wedge}$  (see the proof of Theorem 2.4.1), one can obtain  $\Phi(P \bar{\wedge} Q) = \Phi(P) \bar{\wedge} \Phi(Q)$ , which implies that  $\Phi([P, Q]) = [\Phi(P), \Phi(Q)]$ .

**Definition 2.6.1** *The isomorphism  $\Phi : \mathcal{V} \rightarrow \tilde{\mathcal{V}}$  is called the reciprocal transformation w.r.t.  $\rho$ . Let  $\rho_0$  be the degree zero component of  $\rho$ , if  $\rho = \rho_0$  then  $\Phi$  is called a reciprocal transformation of the first kind, and if  $\rho_0 = 1$  then  $\Phi$  is called a reciprocal transformation of the second kind.*

When restricted to the subspace  $\mathcal{E} \subset \mathcal{V}$  the reciprocal transformation  $\Phi$  has a simple computation formula, we will give it below after some preparations.

We introduce a new coordinate system on  $\mathcal{A}$

$$\tilde{u}^i = u^i, \quad \tilde{u}^{i,s} = \tilde{\partial}^s(\tilde{u}^i).$$

It is easy to see that the transformation  $\{u^{i,s}\} \rightarrow \{\tilde{u}^{i,s}\}$  is invertible, and elements of  $\mathcal{A}$  can be written as differential polynomials in  $\tilde{u}^{i,s}$ . By defining the new derivations  $\tilde{\partial}_{i,s} = \frac{\partial}{\partial \tilde{u}^{i,s}}$ , one can represent  $\tilde{\partial}$  as

$$\tilde{\partial} = \prod_{s \geq 0} \tilde{u}^{i,s+1} \tilde{\partial}_{i,s}.$$

Furthermore, one can define operators  $\tilde{p}_{i,\alpha,s}$ ,  $\tilde{E}_s$  and  $\tilde{E}$  by replacing  $\partial$ ,  $\partial_{i,s}$ ,  $u^{i,s}$  by  $\tilde{\partial}$ ,  $\tilde{\partial}_{i,s}$ ,  $\tilde{u}^{i,s}$  respectively in the original definition of  $p_{i,\alpha,s}$ ,  $E_s$ ,  $E$  given in Section 2.

**Lemma 2.6.2** *Let  $f \in \mathcal{A}$ ,  $\tilde{f} = \rho^{-1}f$ , then the following identities hold true*

$$\delta_i(f) = \rho \tilde{\delta}_i(\tilde{f}) - \prod_{s \geq 0} (-\partial)^s \partial_{i,s}(\rho) \tilde{E}(\tilde{f}), \quad (2.36)$$

$$E(f) = \rho \tilde{E}(\tilde{f}) - \prod_{s \geq 0} \prod_{\alpha \geq 1} u^{i,\alpha} (-\partial)^s \partial_{i,s+\alpha}(\rho) \tilde{E}(\tilde{f}). \quad (2.37)$$

*Proof* We denote the right hand sides of (2.36) and (2.37) by  $g_i$  and  $h$ . According to Corollary 2.1.10, we only need to prove that for any first order differential operator  $D : \mathcal{A} \rightarrow \mathcal{A}$  satisfying  $D\partial = \partial(D - D(1))$ , one has

$$D(f) \sim X^i g_i - a h, \quad (2.38)$$

where  $a = D(1)$  and  $X^i = (D - D(1))(u^i)$ . In fact, such  $D$  corresponds to a quasi-local vector  $X = X^i \theta_i - a \zeta \in \mathcal{E}^1$ , and  $D(f) \sim X(F)$ , where  $F = \int_{\mathbb{R}} f dx$ .

Define  $\tilde{F} = \Phi(F) = \int_{\mathbb{R}} \tilde{f} d\tilde{x}$ , then we have

$$[\Phi(X), \tilde{F}] = \Phi(X)(\tilde{F}) = \Phi_0 X \Phi_0^{-1}(\tilde{F}) = \int_{\mathbb{R}} \rho^{-1} D \rho (\tilde{f}) d\tilde{x}.$$

Denote  $\tilde{D} = \rho^{-1} D \rho$ , one can check that  $\tilde{D}$  is a first order differential operator satisfying  $\tilde{D}\tilde{\partial} = \tilde{\partial} \tilde{D} - \tilde{D}(1)$ , then Lemma 2.1.8 implies

$$\tilde{D}(\tilde{f}) \sim \tilde{X}^i \tilde{\delta}_i(\tilde{f}) - \tilde{a} \tilde{E}(\tilde{f}),$$

where  $\tilde{a} = \tilde{D}(1) = \rho^{-1}D(\rho)$ ,  $\tilde{X}^i = \tilde{D} - \tilde{D}(1)$  ( $\tilde{u}^i = X^i$ ), and

$$A \sim B \Leftrightarrow A - B \in \tilde{\partial}\mathcal{A}.$$

On the other hand,

$$D(f) \sim X(F) = \Phi_0^{-1} [\tilde{\Phi}(X), \tilde{F}] = \Phi_0^{-1} \int \tilde{D}(\tilde{f}) d\tilde{x} = \int \rho \tilde{D}(\tilde{f}) dx,$$

so we have

$$D(f) \sim \rho X^i \tilde{\delta}_i(\tilde{f}) - D(\rho) \tilde{E}(\tilde{f}),$$

then one can obtain (2.38) after integration by parts. The lemma is proved.  $\square$

As we did in the last subsection for Miura type transformations, if we regard  $\delta_i$  and  $E$  as super variables  $\theta_i$  and  $\zeta$  the above lemma gives the transformation rule of super variables under reciprocal transformations. More precisely, we introduce a new family of super variables  $\tilde{\theta}_i^s, \tilde{\zeta}$ , and define the space

$$\tilde{\mathcal{S}} = \mathcal{A} \otimes \wedge^*(\tilde{V}), \text{ where } \tilde{V} = \bigoplus_{i,s} \mathbb{R}\tilde{\theta}_i^s \oplus \mathbb{R}\tilde{\zeta}.$$

We can also define the quotient space  $\tilde{\mathcal{E}} = \tilde{\mathcal{S}}/\tilde{\partial}\tilde{\mathcal{S}}$  which is isomorphic to the space of quasi-local multi-vectors in  $\tilde{\mathcal{V}}$ .

According to Lemma 2.6.2, we introduce the following isomorphism  $\hat{\Phi} : \mathcal{S} \rightarrow \tilde{\mathcal{S}}$  of super commutative algebras

$$\begin{aligned} \hat{\Phi}(f) &= f, \quad f \in \mathcal{A}, \\ \hat{\Phi}(\theta_i^s) &= \rho \tilde{\partial}^s \rho \tilde{\theta}_i - \sum_{t \geq 0} (-\rho \tilde{\partial})^t \partial_{i,t}(\rho) \tilde{\zeta}, \\ \hat{\Phi}(\zeta) &= \rho \tilde{\zeta} - \sum_{s \geq 0} \sum_{\alpha \geq 1} u^{i,\alpha} (-\rho \tilde{\partial})^s \partial_{i,s+\alpha}(\rho) \tilde{\zeta}. \end{aligned}$$

**Theorem 2.6.3** *The restriction of the reciprocal transformation  $\Phi$  on  $\mathcal{E} \subset \mathcal{V}$  is an isomorphism from  $\mathcal{E}$  to  $\tilde{\mathcal{E}}$ . More precisely, let  $\alpha \in \mathcal{S}^p$ , we have*

$$\int \alpha dx = \int \rho^{-1} \hat{\Phi}(\alpha) d\tilde{x}. \quad (2.39)$$

*Proof* The theorem can be easily proved by acting both sides of (2.39) on the local functionals  $\tilde{F}_1, \dots, \tilde{F}_p \in \tilde{\mathcal{F}}$  and using Lemma 2.6.2.  $\square$

**Example 2.6.4** *We consider an evolutionary PDE*

$$\partial_t u^i = X^i, \text{ where } X^i \in \mathcal{A}. \quad (2.40)$$

It corresponds to a local vector  $X = \int^R X^i \theta_i dx \in \mathcal{E}^1$ . Let  $\rho \in \mathcal{A}$  be an invertible element, we can define the reciprocal transformation  $\Phi$  w.r.t.  $\rho$ . From Theorem 2.6.3 it follows that

$$\Phi(X) = \int^Z X^i \tilde{\theta}_i - \rho^{-1} \partial_t(\rho) \tilde{\zeta} d\tilde{x}. \quad (2.41)$$

If  $\rho$  is a conserved density of  $\partial_t$ , i.e. there exists  $\sigma \in \mathcal{A}$  such that  $\partial_t(\rho) = \partial(\sigma)$ , then

$$\Phi(X) = \int^Z X^i - \sigma \tilde{u}^{i,1} \tilde{\theta}_i d\tilde{x}.$$

So we obtain another evolutionary PDE

$$\partial_{\tilde{t}} \tilde{u}^i = X^i - \sigma \tilde{u}^{i,1}. \quad (2.42)$$

This equation coincides with the one that is obtained from (2.40) by the reciprocal transformation (1.6).

If  $\Phi$  is a first kind reciprocal transformation, then the isomorphism  $\hat{\Phi}$  is very simple:

$$\hat{\Phi}(f) = f, \quad \hat{\Phi}(\theta_i^s) = \rho \tilde{\partial}^s \rho \tilde{\theta}_i - \partial_{i,0}(\rho) \tilde{\zeta}, \quad \hat{\Phi}(\zeta) = \rho \tilde{\zeta}. \quad (2.43)$$

When  $\Phi$  is of the second class, we first identify  $\mathcal{E}$  and  $\tilde{\mathcal{E}}$ , and regard  $\Phi$  as an automorphism on  $\mathcal{E}$ , then we have the following formula.

**Theorem 2.6.5** *Let  $\rho = e^f$ , where  $f \in \mathcal{A}_{>0}$ , and let  $\Phi$  be the reciprocal transformation w.r.t.  $\rho$ , then for any  $P \in \mathcal{E}$  we have*

$$\Phi(P) = \exp(\text{ad}_Y)(P),$$

where  $Y = \int^R f \zeta dx \in \mathcal{E}^1$ .

*Proof* The proof of this theorem is similar to that of Theorem 2.5.7, so we omit the details here.  $\square$

**Remark 2.6.6** *Theorem 2.5.7 and Theorem 2.6.5 show that both Miura type transformations of the second kind and reciprocal transformations of the second kind can be expressed as  $\exp(\text{ad}_X)$ , where  $X \in \mathcal{E}_{>0}^1$ . Conversely, by using the Baker-Campbell-Hausdorff formula one can show that every automorphism  $\exp(\text{ad}_X) : \mathcal{E} \rightarrow \mathcal{E}$  with  $X \in \mathcal{E}_{>0}^1$  can be represented as the composition of a Miura type transformation of the second kind and a reciprocal transformation of the second kind. This observation is very important when we study the classification problem of deformations of Jacobi structures (see Section 3.3 below).*

### 3 Jacobi structures

#### 3.1 Definition and examples

A Jacobi structure on a finite dimensional manifold  $M^n$ , as it was introduced by Lichnerowicz in [38], consists of a pair  $(\Lambda, X)$  of a bivector  $\Lambda$  and a vector field  $X$  on  $M$ , they satisfy the conditions

$$[\Lambda, \Lambda] = 2X \wedge \Lambda, \quad [X, \Lambda] = 0. \quad (3.1)$$

It is equivalent to a local Lie algebra structure defined on the space of smooth functions on  $M$  via the following bracket:

$$\{f, g\} = \Lambda(df, dg) + fX(g) - gX(f), \quad \forall f, g \in C^\infty(M).$$

This bracket satisfies the Jacobi identity, and in general (when  $X \neq 0$ ) does not satisfy the Leibniz rule. In this section, we generalize such structures to the infinite jet space of  $M$ , and in this way define Jacobi structures of evolutionary PDEs.

We denote the Lie algebra of derivations on  $\mathcal{A}$  by  $\text{Der}(\mathcal{A})$ , and denote the centralizer of  $\partial$  in  $\text{Der}(\mathcal{A})$  by  $\text{Der}'(\mathcal{A})$ . It is easy to see that the space  $\text{Der}'(\mathcal{A})$  consists of elements of the form

$$D = \sum_{s \geq 0} \partial^s X^i \partial_{i,s}, \quad X^i \in \mathcal{A}.$$

Each such a derivation corresponds to an evolutionary PDE with components in  $\mathcal{A}$ :

$$\partial_t u^i = X^i, \quad X^i \in \mathcal{A}. \quad (3.2)$$

Note that the action of  $D$  on  $f \in \mathcal{A}$  is just  $\partial_t f$ .

Since elements of  $\text{Der}'(\mathcal{A})$  commute with  $\partial$ ,  $\mathcal{F}$  has a natural  $\text{Der}'(\mathcal{A})$ -module structure given by

$$\pi : \text{Der}'(\mathcal{A}) \rightarrow \mathcal{V}^1 = \text{End}(\mathcal{F}, \mathcal{F}), \quad \pi(D) \begin{matrix} \int \\ f dx \end{matrix} = \begin{matrix} \int \\ D(f) dx \end{matrix}.$$

The image of  $\pi$  is just  $\mathcal{E}_{loc}^1$  (since  $D(f) \sim X^i \delta_i(f)$ ), and it follows from Lemma 2.1.7 that the kernel of  $\pi$  is  $\mathbb{R} \partial$ . So we have the following isomorphism:

$$\mathcal{E}_{loc}^1 \cong \text{Der}'(\mathcal{A}) / \mathbb{R} \partial.$$

This quotient has a nice physical explanation. Given an evolutionary PDE (3.2), we can convert it to

$$\partial_t u^i = X^i + c u^{i,1}$$

by performing the following Galilean transformation

$$x \mapsto x - ct, \quad t \mapsto t,$$

so the space  $\mathcal{E}_{loc}^1$  is just the space of equivalence classes of evolutionary PDEs modulo Galilean transformations. In this paper, we consider these equivalence classes only, and call elements of  $\mathcal{E}_{loc}^1$  evolutionary PDEs for short.

**Definition 3.1.1** A quasi-local bivector  $P \in \mathcal{E}^2$  is called a Jacobi structure if  $[P, P] = 0$ . An evolutionary PDE  $X \in \mathcal{E}_{loc}^1$  is said to possess a Jacobi structure if there exists a Jacobi structure  $P$  and a local functional  $F \in \mathcal{F}$  such that  $X = [P, F]$ . Here  $F$  is called the Hamiltonian.

By using integration by parts one can always represent a Jacobi structure in the following form:

$$P = \int \frac{1}{2} \theta_i \alpha_s^{ij} \partial^s \theta_j + \zeta X^i \theta_i \, dx,$$

where  $\alpha_s^{ij}, X^i \in \mathcal{A}$ , and the matrix differential operator  $\alpha = (\alpha_s^{ij} \partial^s)$  satisfies the skew-symmetry condition

$$\alpha + \alpha^\dagger = 0 \tag{3.3}$$

with  $\alpha^\dagger = (((-\partial)^s \alpha_s^{ji}))$ . The bivector  $P_0 = \frac{1}{2} \theta_i \alpha^{ij} \theta_j \in \mathcal{S}_{loc}^2$  is called the local part of  $P$  and  $X = X^i \theta_i \in \mathcal{S}_{loc}^1$  is called, following the notation of [25], the structure flow of  $P$ . In what follows a Jacobi structure  $P$  is often represented as  $P = (P_0, X)$ , or  $P \sim P_0 + \zeta X$ .

**Lemma 3.1.2** Let  $X \in \mathcal{E}_{loc}^1, P \in \mathcal{E}^2$  be a Jacobi structure of  $X$ , then the structure flow of  $P$  is a symmetry of  $X$ .

*Proof* By definition, there exists  $F \in \mathcal{F}$  such that  $X = [P, F]$ , so we have  $[X, P] = 0$ . Note that the derivation  $\partial_\zeta : \mathcal{S} \rightarrow \mathcal{S}$  satisfies  $[\partial_\zeta, \partial] = 0$ , so it induces a map  $\partial_\zeta : \mathcal{E} \rightarrow \mathcal{E}$ . By the action of  $\partial_\zeta$ , one can obtain

$$0 = \partial_\zeta ([X, P]) = [X, \partial_\zeta(P)],$$

where  $\partial_\zeta(P)$  is just the structure flow of  $P$ . The lemma is proved.  $\square$

**Lemma 3.1.3** Let  $P = (P_0, X)$  be a Jacobi structure,  $H \in \mathcal{F}$  be a local functional, then  $[P, H]$  is local if and only if  $[X, H] = 0$ .

*Proof* The locality of  $[P, H]$  means that  $\partial_\zeta[P, H] = 0$ , so we have

$$\begin{aligned} 0 &= \partial_\zeta[P, H] \sim \partial_\zeta \left( \delta^i(P) \delta_i(H) + \partial_\zeta(P) \hat{E}(H) \right) \\ &= \partial_\zeta \left( \delta^i(\zeta X) \delta_i(H) \right) \sim -[X, H], \end{aligned}$$

the lemma is proved.  $\square$

The above lemma means that for a given Jacobi structure  $P = (P_0, X)$ , the admissible Hamiltonians must be conserved quantities of the structure flow  $X$ . This is quite different from the local case.

**Example 3.1.4** *The simplest Jacobi structures are the ones with degree zero. Let  $P = (P_0, X) \in \mathcal{E}^2$  be a Jacobi structure such that  $\deg(P) = 0$ , then  $P_0$  and  $X$  are just a bivector and a vector field on  $M$  respectively. The condition  $[P, P] = 0$  implies*

$$[P_0, P_0] + 2X P_0 = 0, \quad [P_0, X] = 0,$$

so the pair  $(P_0, -X)$  gives a classical Jacobi structure on  $M$  [38].

Let  $P \sim \frac{1}{2}\alpha^{ij}\theta_i\theta_j + \zeta X^i\theta_i$  be a Jacobi structure of degree zero, and  $\rho$  be a nowhere zero smooth function on  $M$ . One can define the reciprocal transformation  $\Phi$  w.r.t.  $\rho$ . The isomorphism  $\hat{\Phi}$  is given by (2.43), so we obtain the reciprocal transformation of  $P$

$$\Phi(P) \sim \frac{1}{2}\tilde{\alpha}^{ij}\tilde{\theta}_i\tilde{\theta}_j + \tilde{\zeta}\tilde{X}^i\tilde{\theta}_i,$$

where  $\tilde{\alpha}^{ij} = \rho\alpha^{ij}$ ,  $\tilde{X}^i = \rho X^i + \alpha^{ij}\partial_{j,0}(\rho)$ . This is in fact a conformal change of classical Jacobi structures [10, 30].

The Jacobi structures of degree 1 are more interesting for us. Let  $P = (P_0, X)$  be a Jacobi structure where

$$P_0 = \frac{1}{2}g^{ij}(u)\theta_i\theta_j^1 + \Gamma_k^{ij}(u)u^{k,1}\theta_i\theta_j, \quad X = V_k^i(u)u^{k,1}\theta_i, \quad (3.4)$$

and we assume that  $\det(g^{ij}) \neq 0$ . The skew-symmetry condition (3.3) is equivalent to

$$g^{ij} = g^{ji}, \quad \Gamma_k^{ij} + \Gamma_k^{ji} = g_{(k,0)}^{ij} = \frac{\partial g^{ij}}{\partial u^k},$$

so  $g^{ij}$  is a contravariant metric on  $M$ , we denote its inverse by  $g_{ij}$ , and define  $\Gamma_{kl}^j = -g_{ki}\Gamma_l^{ij}$ ,  $V_{kj} = g_{ki}V_j^i$ .

The following theorem was first proved by Ferapontov in [23], here we give another proof to illustrate the usage of our formula of the Schouten-Nijenhuis bracket (2.23).

**Theorem 3.1.5** *A bivector  $P = (P_0, X)$  of the form (3.4) is a Jacobi structure if and only if the following conditions are satisfied*

$$\begin{aligned} \Gamma_{kl}^j &= \Gamma_{lk}^j, \quad V_{kj} = V_{jk}, \quad \nabla_k V_{lj} = \nabla_l V_{kj}, \\ R_{ijkl} &= g_{ik}V_{jl} + g_{jl}V_{ik} - g_{jk}V_{il} - g_{il}V_{jk}, \end{aligned} \quad (3.5)$$

where  $\nabla$  and  $R_{ijkl}$  are the Levi-Civita connection and Riemannian curvature tensor of  $g_{ij}$  respectively.

*Proof* We need to check that the condition  $[P, P] = 0$  is equivalent to the conditions listed in the theorem. From (2.23) it follows that

$$\frac{1}{2}[P, P] \sim \delta^i(P_0 + \zeta X)\delta_i(P_0 + \zeta X) + X\hat{E}(P_0 + \zeta X).$$

By using the following identities,

$$\begin{aligned}\delta^i(\zeta X) &= -\zeta \delta^i(X), \quad \delta_i(\zeta X) = \zeta \delta_i(X) - u^{k,1}\theta_k \quad V_i^l\theta_l, \\ \hat{E}(\zeta X) &= \zeta X, \quad \hat{E}(P_0) = 2P_0,\end{aligned}$$

one can obtain  $\frac{1}{2}[P, P] \sim A - \zeta B$ , where

$$\begin{aligned}A &= \frac{1}{2}[P_0, P_0]_{pr} - \delta^i(P_0) \quad u^{k,1}\theta_k \quad V_i^l\theta_l + 2X P_0, \\ B &= [P_0, X]_{pr} - \delta^i(X) \quad u^{k,1}\theta_k \quad V_i^l\theta_l.\end{aligned}$$

The condition  $A - \zeta B \sim 0$  implies  $B \sim 0$ . Note that  $B \in \mathcal{S}_{loc}^2$ , so we have  $\delta^i(B) = 0$  (see the proof of Lemma 2.4.10). We rewrite  $B$  in the following form

$$B = E^{ij}\theta_i^1\theta_j^1 + F_k^{ij}u^{k,1}\theta_i\theta_j^1 + H_{kl}^{ij}u^{k,1}u^{l,1}\theta_i\theta_j,$$

then we have

$$0 = \partial_2^i\delta^j(B) = \partial_2^i \quad \partial_0^j - \partial\partial_1^j \quad (B) = \partial_1^j\partial_1^i(B) = E^{ij} - E^{ji}, \quad (3.6)$$

$$0 = \partial_{k,2}\delta^j(B) = \partial_{k,2} \quad \partial_0^j - \partial\partial_1^j \quad (B) = -\partial_{k,1}\partial_1^j(B) = F_k^{ij}\theta_i, \quad (3.7)$$

so we have  $\delta^i(B) = H_{kl}^{ij} - H_{kl}^{ji} \quad u^{k,1}u^{l,1}\theta_j = 0$ , which implies

$$H_{kl}^{ij} - H_{kl}^{ji} + H_{lk}^{ij} - H_{lk}^{ji} = 0. \quad (3.8)$$

The conditions (3.6), (3.7), (3.8) show that  $B = 0$ , so we have  $A \sim 0$  and  $\delta^i(A) = 0$ . Then by straightforward computation, one can obtain  $\Gamma_{kl}^j = \Gamma_{lk}^j$  from  $\partial_2^i\delta^j(A) = 0$ , and  $V_{kj} = V_{jk}$  from (3.6), and  $\nabla_k V_{lj} = \nabla_l V_{kj}$  from (3.7), and  $R_{ijkl}$  from  $\partial_{k,2}\delta^j(A) = 0$ . So we proved that the four conditions are necessary.

The sufficiency part can be proved by a straightforward but lengthy computation, we omit it here.  $\square$

**Definition 3.1.6** *A bivector  $P \sim P_0 + \zeta X \in \mathcal{E}^2$  is called a Jacobi structure of hydrodynamic type if  $(P_0, X)$  is given by (3.4) and  $\det(g^{ij}) \neq 0$ .*

In what follows, we will study reciprocal transformations of hydrodynamic Jacobi structures. Note that reciprocal transformations preserve the Schouten-Nijenhuis bracket, so the reciprocal transformation of a Jacobi structure is still a Jacobi structure. The following lemma that is given in [2, 25] is very useful when studying reciprocal transformations of hydrodynamic Jacobi structures. It can be proved by using the relation (2.43) and our general transformation formula (2.39).

**Lemma 3.1.7 ([2, 25])** *Let  $P = (P_0, X)$  be a Jacobi structure of hydrodynamic type and  $\rho$  be a nowhere zero smooth function on  $M$ . Denote the reciprocal transformation w.r.t.  $\rho$  by  $\Phi$ , then  $\Phi(P) = (\tilde{P}_0, \tilde{X})$  has the expression*

$$\tilde{P}_0 = \frac{1}{2} \quad \tilde{g}^{ij}(u)\tilde{\theta}_i\tilde{\theta}_j^1 + \tilde{\Gamma}_k^{ij}(u)\tilde{u}^{k,1}\tilde{\theta}_i\tilde{\theta}_j, \quad \tilde{X} = \tilde{V}_k^i(u)\tilde{u}^{k,1}\tilde{\theta}_i, \quad (3.9)$$

where

$$\tilde{g}^{ij} = \rho^2 g^{ij}, \quad \tilde{V}_k^i = \rho^2 V_k^i + \rho \nabla^i \nabla_k \rho - \frac{1}{2} (\nabla^l \rho) (\nabla_l \rho) \delta_k^i \quad (3.10)$$

and  $\nabla$  is the Levi-Civita connection of the metric  $g_{ij}$ .

*Proof of Theorem 1.1* According to the results of Dubrovin and Novikov [18, 19], we only need to show that there exists a reciprocal transformation  $\Phi$  such that  $\Phi(P)$  is local. Let the Jacobi structure of hydrodynamic type  $P = (P_0, X)$  be given by (3.4). From the condition (3.5), one can obtain the Ricci and scalar curvatures of  $g_{ij}$

$$R_{ij}(g) = (n-2)V_{ij} + V_k^k g_{ij}, \quad R(g) = 2(n-1)V_k^k$$

which imply that both the Weyl tensor and the Cotton tensor of  $g_{ij}$  vanish, so  $g_{ij}$  is a conformally flat metric.

Let  $g_{ij} = \rho^{-2} \eta_{ij}$ , where  $\eta_{ij}$  is a flat metric, and  $\rho^{-2}$  is the conformal factor. Note that  $\rho^{-1}$  is nowhere zero on  $M$ , so we can define a reciprocal transformation  $\Phi$  w.r.t.  $\rho^{-1}$ . Let  $\Phi(P) = Q = (Q_0, Y)$ , then it is easy to see that the metric of  $Q_0$  is just the flat metric  $\eta_{ij}$ . So without loss of generality we can assume that  $g_{ij}$  has been converted to a flat metric  $\eta_{ij}$ .

When  $n \geq 3$ , one obtain

$$V_{ij} = \frac{R_{ij}(\eta)}{n-2} - \frac{g_{ij}R(\eta)}{2(n-1)(n-2)} = 0,$$

so the theorem has been proved in this case.

Now we assume  $n = 2$ . In this case, the tensor  $V_{ij}$  is not determined by the curvatures, so it may not be zero. We choose  $u^1, u^2$  to be the flat coordinates of metric  $\eta$ , then  $\eta_{ij}$  becomes a constant symmetric matrix. There are two possible cases up to linear transformations

$$(\eta_{ij}) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{or} \quad (\eta_{ij}) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

We only consider the first case since the second one is similar and easier.

We denote the components of the tensor  $V_{ij}$  by

$$a = V_{11}, \quad b = V_{12} = V_{21}, \quad c = V_{22},$$

and  $u^1 = x, u^2 = y$ , then the conditions (3.5) imply

$$a + c = 0, \quad a_y = b_x, \quad b_y = c_x.$$

So if we introduce  $z = x + y\sqrt{-1}$ ,  $f = a - b\sqrt{-1}$ , then  $f$  is holomorphic in  $z$ .

On the other hand, if there exists  $\rho$  such that  $\Phi(P)$  is local, then this  $\rho$  must satisfy the following equation which is implied by the formula (3.10):

$$\rho^2 V_{ij} + \rho \nabla_i \nabla_j \rho - \frac{1}{2} (\nabla^l \rho) (\nabla_l \rho) \eta_{ij} = 0.$$

The components of the above equation read

$$\frac{\rho_x^2 + \rho_y^2}{2\rho^2} - \frac{\rho_{xx}}{\rho} = a, \quad -\frac{\rho_{xy}}{\rho} = b, \quad \frac{\rho_x^2 + \rho_y^2}{2\rho^2} - \frac{\rho_{yy}}{\rho} = -a. \quad (3.11)$$

We introduce a new function

$$\chi = -\frac{\rho_x}{\rho} + \frac{\rho_y}{\rho} \sqrt{-1},$$

then the equations (3.11) imply that  $\chi$  is holomorphic in  $z$  and satisfies the following Riccati equation

$$\chi_z - \frac{\chi^2}{2} = f.$$

Since  $f$  is holomorphic, this Riccati equation has solutions. So we can find a reciprocal transformation  $\Phi$  such that  $\Phi(P)$  is local. The  $n = 2$  case is proved.

The  $n = 1$  case is similar to the  $n = 2$  case, we omit the details here.  $\square$

### 3.2 Locality problem

In this subsection, we investigate the following problem: *Under what conditions is the reciprocal transformation of a Jacobi structure local, i.e. a Hamiltonian structure?*

We fix an invertible  $\rho \in \mathcal{A}$ , and denote by  $\Phi$  the reciprocal transformation w.r.t.  $\rho$ . The following local functionals are very useful in this subsection.

$$\Lambda = \int \rho dx \in \mathcal{F}, \quad \tilde{\Lambda} = \int 1 d\tilde{x} \in \tilde{\mathcal{F}}.$$

It is easy to see that  $\Phi(\Lambda) = \tilde{\Lambda}$ ,  $[\tilde{P}, \tilde{\Lambda}] = -\partial_{\tilde{\zeta}}(\tilde{P})$ .

**Lemma 3.2.1** *Let  $\tilde{P} \in \tilde{\mathcal{E}}^p$  with  $p \neq 2$ , then  $\tilde{P}$  is local if and only if  $\partial_{\tilde{\zeta}}(\tilde{P}) = 0$ .*

*Proof* We take a representative of  $\tilde{P} = \int \tilde{P}_0 + \tilde{\zeta} \tilde{X} d\tilde{x}$ , where  $\tilde{P}_0 \in \tilde{\mathcal{S}}_{loc}^p$ ,  $\tilde{X} \in \tilde{\mathcal{S}}_{loc}^{p-1}$ . The condition  $\partial_{\tilde{\zeta}}(\tilde{P}) = \int \tilde{X} d\tilde{x} = 0$  implies that there exists  $\tilde{Y} \in \tilde{\mathcal{S}}^{p-1}$  such that  $\tilde{X} = \tilde{\partial}(\tilde{Y})$ . Suppose  $\tilde{Y} = \tilde{Y}_0 + \tilde{Y}_1 \tilde{\zeta}$  with  $\tilde{Y}_0, \tilde{Y}_1$  local, then the condition  $\tilde{X} \in \tilde{\mathcal{S}}_{loc}^{p-1}$  implies that  $\tilde{Y}_1$  must be a constant. But we have assume  $p \neq 2$ ,  $\tilde{Y}_1 \tilde{\zeta}$  cannot be element of  $\tilde{\mathcal{S}}^{p-1}$  unless  $\tilde{Y}_1 = 0$ , so  $\tilde{Y}$  is also local, then

$$\tilde{P} \sim \tilde{P}_0 + \tilde{\zeta} \tilde{X} = \tilde{P}_0 + \tilde{\zeta} \tilde{\partial}(\tilde{Y}) \sim \tilde{P}_0 - \tilde{\partial}(\tilde{\zeta}) \tilde{Y},$$

which is local. The lemma is proved.  $\square$

**Theorem 3.2.2** *Let  $P \in \mathcal{E}^p$  with  $p \neq 2$ , then  $\Phi(P)$  is local if and only if  $[P, \Lambda] = 0$ .*

*Proof* According to the above lemma,  $\Phi(P)$  is local if and only if  $\partial_{\tilde{\zeta}}(\Phi(P)) = 0$ , which is equivalent to  $[\Phi(P), \tilde{\Lambda}] = 0$ , but we have  $[\Phi(P), \tilde{\Lambda}] = \Phi([P, \Lambda])$ , so the theorem is proved.  $\square$

When  $p = 2$ , it is easy to see that the condition  $[P, \Lambda] = 0$  is still necessary to imply the locality of  $\Phi(P)$ , but not sufficient. According to the proof of Lemma 3.2.1, we have the following definition.

**Definition 3.2.3** *Let  $P \in \mathcal{E}^2$ , and  $\rho$  be an invertible differential polynomial satisfying  $[P, \Lambda] = 0$ , then the reciprocal transformation  $\Phi(P)$  must take the following form:*

$$\Phi(P) = \int \tilde{P}_0 + z \tilde{u}^{i,1} \tilde{\theta}_i \tilde{\zeta} \, d\tilde{x},$$

where  $\tilde{P}_0 \in \tilde{\mathcal{S}}_{loc}^2$  and  $z \in \mathbb{R}$ . The constant  $z$  is called the nonlocal charge of the pair  $(P, \rho)$ , and is denoted by  $z(P, \rho)$ .

**Theorem 3.2.4** *Let  $P \in \mathcal{E}^2$ , then  $\Phi(P)$  is local if and only if  $[P, \Lambda] = 0$ ,  $z(P, \rho) = 0$ .*

**Remark 3.2.5** *When  $P$  is a Jacobi structure of hydrodynamic type, the above result was proved by Abenda and Grave. They also considered locality problems with respect to more general reciprocal transformations of hydrodynamic Hamiltonian structures or bihamiltonian structures, see [2, 3].*

In the remaining part of this subsection, we will give a computation formula for the nonlocal charge of a Jacobi structure with a hydrodynamic leading term.

Let  $P = (P_0, X)$  be a Jacobi structure, and  $\rho$  be an invertible differential polynomial satisfying  $[P, \Lambda] = 0$ , then Lemma 3.1.3 shows that  $[X, \Lambda] = 0$ , so there exists  $\sigma \in \mathcal{A}$  such that

$$\partial^s(\delta^i(X))\partial_{i,s}(\rho) = \partial(\sigma).$$

Suppose the leading term of  $P$  is of hydrodynamic type,

$$P_0 = \frac{1}{2} g^{ij} \theta_i \theta_j + \Gamma_k^{ij} u^{k,1} \theta_i \theta_j + \dots, \quad X = V_k^i u^{k,1} + \dots \theta_i,$$

and suppose the decomposition of  $\rho$  and  $\sigma$  w.r.t. the gradation (2.2) reads

$$\rho = \rho_0 + \dots, \quad \sigma = \sigma_0 + \dots,$$

here  $\dots$  denote the terms with higher degrees, then it is easy to see

$$V_k^i \nabla_i(\rho_0) = \nabla_k(\sigma_0). \tag{3.12}$$

By straightforward computation, one can obtain

$$[P, \Lambda] = - \int \nabla^i \nabla_k(\rho_0) + \rho_0 V_k^i + \sigma_0 \delta_k^i u^{k,1} + \dots \theta_i \, dx,$$

if  $[P, \Lambda] = 0$ , then there exists  $c \in \mathbb{R}$  such that

$$\nabla^i \nabla_k (\rho_0) + \rho_0 V_k^i + \sigma_0 \delta_k^i = c \delta_k^i. \quad (3.13)$$

On the other hand, Lemma 3.1.7 shows that the nonlocal part of  $\Phi(P)$  is

$$(\Phi(P))_{nl} = \int \tilde{\zeta} \tilde{V}_k^i \tilde{u}^{k,1} \tilde{\theta}_i d\tilde{x},$$

where  $\tilde{V}_k^i$  reads

$$\begin{aligned} \tilde{V}_k^i &= \rho_0^2 V_k^i + \rho_0 \nabla^i \nabla_k \rho_0 - \frac{1}{2} (\nabla^l \rho_0) (\nabla_l \rho_0) \delta_k^i \\ &= \rho_0 (c - \sigma_0) - \frac{1}{2} (\nabla^l \rho_0) (\nabla_l \rho_0) \delta_k^i, \end{aligned}$$

so we have

$$(\Phi(P))_{nl} = \int \frac{1}{2} (\nabla^l \rho_0) (\nabla_l \rho_0) + \rho_0 (\sigma_0 - c) \tilde{u}^{i,1} \tilde{\theta}_i \tilde{\zeta} d\tilde{x},$$

comparing with Definition 3.2.3, we obtain

$$z(P, \rho) = \frac{1}{2} (\nabla^l \rho_0) (\nabla_l \rho_0) + \rho_0 (\sigma_0 - c). \quad (3.14)$$

By using the equations (3.12) and (3.13), one can easily show that the right hand side of (3.14) is indeed a constant, and is independent of the choice of  $\sigma$ , so we get the correct formula for the nonlocal charge of  $(P, \rho)$ .

**Corollary 3.2.6** *Let  $P$  be a quasi-local bivector whose leading term is a Jacobi structure of hydrodynamic type, suppose  $\rho, \rho'$  are two invertible differential polynomials satisfying  $\rho \sim \rho'$ ,  $\Phi, \Phi'$  are the reciprocal transformations w.r.t.  $\rho, \rho'$  respectively, then  $\Phi(P)$  is local if and only if  $\Phi'(P)$  is local.*

*Proof* According to Theorem 3.2.4,  $\Phi(\mathbb{R})$  is local if and only if  $[P, \int \rho dx] = 0$  and  $z(P, \rho) = 0$ . Since  $\rho \sim \rho'$ , we have  $[P, \int \rho' dx] = 0$ . Note that  $z(P, \rho)$  (3.14) only depends on the degree zero part of  $\rho$ , which coincides with the one of  $\rho'$ , so we also have  $z(P, \rho') = 0$ . Then by using Theorem 3.2.4 again, we obtain that  $\Phi'(P)$  is local. The corollary is proved.  $\square$

### 3.3 Deformation and cohomology

In Section 3.1, we proved that every Jacobi structure of hydrodynamic type is equivalent to a Hamiltonian structure of hydrodynamic type modulo reciprocal transformations.

Then by using the classical result of Dubrovin and Novikov [18, 19] that every Hamiltonian structure of hydrodynamic type is equivalent to a Hamiltonian structure of the form

$$P_{norm} = \int \frac{1}{2} \eta^{ij} \theta_i \theta_j^1 dx \quad (3.15)$$

modulo coordinate transformations, we arrive at the classification of Jacobi structures of hydrodynamic type: *every Jacobi structure of hydrodynamic type is equivalent to a Hamiltonian structure of the form  $P_{norm}$ .*

In this subsection, we will generalize the above classification result to more general Jacobi structures.

Let  $P$  be an arbitrary Jacobi structure. Note that  $(\mathcal{E}, d_P)$  forms a differential graded Lie algebra (DGLA), where  $d_P = \text{ad}_P$  is the adjoint action of  $P$ :

$$\text{ad}_P : \mathcal{E} \rightarrow \mathcal{E}, \quad Q \mapsto \text{ad}_P(Q) = [P, Q],$$

so we immediately obtain the following definitions and results along the line of the general philosophy about the relationship between deformation problems and the cohomologies of the associated DGLAs.

**Definition 3.3.1** *Let  $P_0 \in \mathcal{E}^2$  be a Jacobi structure, we say  $P = P_0 + Q \in \mathcal{E}^2$  is a deformation of  $P_0$  if  $P$  is also a Jacobi structure and  $\nu(Q) > \nu(P_0)$ . Let  $P, P'$  be two deformations of  $P_0$ , we say  $P$  is equivalent to  $P'$  if there exists  $X \in \mathcal{E}^1$  such that  $\nu(X) > 0$  and  $e^{\text{ad}_X}(P) = P'$ . A deformation  $P$  is called trivial if it is equivalent to  $P_0$ .*

Note that in the above definition the function  $\nu$  is introduced in Definition 2.5.5.

**Definition 3.3.2** *Let  $P \in \mathcal{E}^2$  be a Jacobi structure,  $Q \in \mathcal{E}^2$  is called an infinitesimal deformation of  $P$  if  $\nu(Q) > \nu(P)$  and  $d_P(Q) = 0$ . Let  $Q, Q'$  be two infinitesimal deformations of  $P$ , they are called equivalent if there exists  $X \in \mathcal{E}^1$  such that  $\nu(X) > 0$  and  $Q' - Q = d_P(X)$ . An infinitesimal deformation  $Q$  is called trivial if it is equivalent to 0.*

**Definition 3.3.3** *Let  $P \in \mathcal{E}^2$  be a Jacobi structure, due to  $d_P^2 = 0$  we have the following complex*

$$0 \longrightarrow \mathcal{E}^0 \xrightarrow{d_P} \mathcal{E}^1 \xrightarrow{d_P} \mathcal{E}^2 \xrightarrow{d_P} \dots,$$

*its cohomology is called the Lichnerowicz-Jacobi cohomology of  $P$*

$$H^i(\mathcal{E}, P) = \frac{\text{Ker}(d_P : \mathcal{E}^i \rightarrow \mathcal{E}^{i+1})}{\text{Im}(d_P : \mathcal{E}^{i-1} \rightarrow \mathcal{E}^i)}. \quad (3.16)$$

If  $P$  is homogeneous, i.e.  $P = P_{\nu(P)}$ , then  $H^i(\mathcal{E}, P)$  possesses a gradation

$$H^i(\mathcal{E}, P) = \bigvee_{d \geq 0} H_d^i(\mathcal{E}, P), \quad H_d^i(\mathcal{E}, P) = \frac{\text{Ker } d_P : \mathcal{E}_d^i \rightarrow \mathcal{E}_{d+\nu(P)}^{i+1}}{\text{Im } d_P : \mathcal{E}_{d-\nu(P)}^{i-1} \rightarrow \mathcal{E}_d^i},$$

where  $\mathcal{E}_d^i = \{f \in \mathcal{E}^i \mid \deg f = d\}$ .

**Remark 3.3.4** Let  $\Phi$  be a reciprocal transformation, and  $P$  be a Jacobi structure. Since  $\Phi$  preserves the Schouten-Nijenhuis bracket, it induces an isomorphism of complexes  $\Phi : (\mathcal{E}, d_P) \rightarrow (\tilde{\mathcal{E}}, d_{\Phi(P)})$ , which implies

$$H^*(\mathcal{E}, P) \cong H^*(\tilde{\mathcal{E}}, \Phi(P)),$$

so reciprocal transformations preserve the Lichnerowicz-Jacobi cohomology. When  $P$  is a Jacobi structure of degree zero (see Example 3.1.4), then  $H_0^*(\mathcal{E}, P)$  coincides with the classical Lichnerowicz-Jacobi cohomology defined in [36]. If  $\Phi$  is given by a nowhere zero smooth function on  $M$ , then  $\Phi(P)$  is just a conformal change of  $P$  (see Example 3.1.4), so the above isomorphism shows that the Lichnerowicz-Jacobi cohomology is invariant under conformal changes of the Jacobi structure. This fact was first shown in [36], our present proof seems simpler than the one given there.

**Proposition 3.3.5** Let  $P$  be a homogeneous Jacobi structure.

a) The space of equivalence classes of infinitesimal deformations of  $P$  is given by  $H_{>\nu(P)}^2(\mathcal{E}, P)$ . In particular, every deformation of  $P$  is trivial if and only if  $H_{>\nu(P)}^2(\mathcal{E}, P)$  vanishes.

b) If  $H_{>2\nu(P)}^3(\mathcal{E}, P)$  vanishes, then every infinitesimal deformation can be extended to a genuine deformation, and the space of equivalence classes of deformations of  $P$  is just  $H_{>\nu(P)}^2(\mathcal{E}, P)$ .

In the next subsection, we will prove the following theorem.

**Theorem 3.3.6** Let  $P$  be a Jacobi structure of hydrodynamic type, then we have

$$H_{>0}^q(\mathcal{E}, P) \cong 0, \text{ for } q = 0, 1, 2, \dots$$

From this theorem we see that any deformation of a Jacobi structure of hydrodynamic type can be transformed to  $P_{norm}$  by a series of Miura type transformations and reciprocal transformations. So the Main Theorem is proved.

**Corollary 3.3.7** Let  $P$  be a deformation of a Jacobi structure of hydrodynamic type, if an  $X \in \mathcal{E}_{loc}^1$  satisfies  $[P, X] = 0$ , then the associated system of evolutionary PDEs has a Jacobi structure, i.e. there exists  $H \in \mathcal{F}$  such that  $X = [P, H]$ .

### 3.4 Computation of cohomologies

We fix a Jacobi structure of hydrodynamic type  $P \in \mathcal{E}^2$  in this subsection. Note that Miura type transformations and reciprocal transformations preserve the Lichnerowicz-Jacobi cohomologies, so we can assume  $P = \frac{1}{2}\eta^{ij}\theta_i\theta_j dx$  without loss of generality.

To every  $Q \in \mathcal{E}$ , one can associate a first order differential operator  $D_Q : \mathcal{S} \rightarrow \mathcal{S}$  (see (2.25)) such that

$$D_Q(\alpha) dx = [Q, \alpha dx], \text{ for all } \alpha \in \mathcal{S}.$$

For our fixed  $P$ , the operator  $D_P$  reads

$$D_P = \eta^{ij} \theta_i \theta_j^1 \partial_\zeta + \eta^{ij} \theta_j^{s+1} \partial_{i,s}$$

and one can prove the following identities

$$D_P^2 = 0, \quad [D_P, \partial] = 0. \quad (3.17)$$

Thus we obtain a complex  $(\mathcal{S}, D_P)$

$$0 \longrightarrow \mathcal{S}^0 \xrightarrow{D_P} \mathcal{S}^1 \xrightarrow{D_P} \mathcal{S}^2 \xrightarrow{D_P} \dots,$$

and a short exact sequence of complexes

$$0 \longrightarrow (\mathcal{S}, D_P) \xrightarrow{\partial} (\mathcal{S}, D_P) \xrightarrow{J} (\mathcal{E}, d_P) \longrightarrow 0.$$

Then by using the long exact sequence theorem, we obtain the following lemma.

**Lemma 3.4.1** *For every  $d \geq 0$ , we have the long exact sequence:*

$$\dots \longrightarrow H_{d-1}^p(\mathcal{S}) \xrightarrow{\partial^*} H_d^p(\mathcal{S}) \xrightarrow{J^*} H_d^p(\mathcal{E}) \longrightarrow H_d^{p+1}(\mathcal{S}) \longrightarrow \dots \quad (3.18)$$

where  $H_d^p(\mathcal{S}) = H_d^p(\mathcal{S}, D_P)$ ,  $H_d^p(\mathcal{E}) = H_d^p(\mathcal{E}, d_P)$ .

The complex  $(\mathcal{S}, D_P)$  has a subcomplex  $(\mathcal{S}_{loc}, D'_P)$ , where  $D'_P$  is the restriction of  $D_P$  on  $\mathcal{S}_{loc}$ :

$$D'_P = \eta^{ij} \theta_j^{s+1} \partial_{i,s}.$$

The operator  $D'_P$  satisfies  $D'_P \partial_\zeta + \partial_\zeta D'_P = 0$ , so we have another short exact sequence of complexes

$$0 \longrightarrow (\mathcal{S}_{loc}, D'_P) \xrightarrow{i} (\mathcal{S}, D_P) \xrightarrow{\partial_\zeta} (\mathcal{S}_{loc}, -D'_P) \longrightarrow 0,$$

and the following lemma.

**Lemma 3.4.2** *For every  $d \geq 0$ , we have the long exact sequence:*

$$\dots \longrightarrow H_d^p(\mathcal{S}_{loc}) \xrightarrow{i^*} H_d^p(\mathcal{S}) \xrightarrow{\partial_\zeta^*} H_d^{p-1}(\mathcal{S}_{loc}) \longrightarrow H_{d+1}^{p+1}(\mathcal{S}_{loc}) \longrightarrow \dots \quad (3.19)$$

where  $H_d^p(\mathcal{S}_{loc}) = H_d^p(\mathcal{S}_{loc}, D'_P) \cong H_d^p(\mathcal{S}_{loc}, -D'_P)$ .

Recall that the algebra  $\mathcal{S}_{loc}$  is defined as

$$\mathcal{S}_{loc} = \mathcal{A} \otimes \wedge^*(V_{loc}), \quad \text{where } V_{loc} = \bigoplus_{i,s} (\mathbb{R}\theta_i^s).$$

We denote  $\theta^{i,s} = \eta^{ij}\theta_j^s$ , and decompose  $V_{loc}$  as  $V_{loc} = V_{loc}^0 \oplus V_{loc}^{>0}$ , where

$$V_{loc}^0 = \bigvee_{i=1}^{\mathbb{N}} \mathbb{R}\theta^{i,0} \cong \mathbb{R}^n, \quad V_{loc}^{>0} = \bigvee_{i=1}^{\mathbb{N}} \bigvee_{s>0} \mathbb{R}\theta^{i,s},$$

then  $\mathcal{S}_{loc}$  has the following tensor product decomposition

$$\mathcal{S}_{loc} = \mathcal{S}_{loc}^{>0} \otimes \wedge^*(V_{loc}^0), \quad \text{where } \mathcal{S}_{loc}^{>0} = \mathcal{A} \otimes \wedge^*(V_{loc}^{>0}),$$

and the differential  $D'_P$  becomes  $D'_P = \theta^{i,s+1}\partial_{i,s}$ , which is a differential on  $\mathcal{S}_{loc}^{>0}$ . By using the universal coefficient theorem, we obtain the following lemma.

**Lemma 3.4.3**

$$H^*(\mathcal{S}_{loc}, D'_P) \cong H^*(\mathcal{S}_{loc}^{>0}, D'_P) \otimes \wedge^*(V_{loc}^0). \quad (3.20)$$

By regarding  $\theta^{i,s+1}$  as  $du^{i,s}$ , it is easy to see that the complex  $(\mathcal{S}_{loc}^{>0}, D'_P)$  is just the de Rham complex  $(J^\infty(M), d_{dR})$  of  $J^\infty(M)$  with coefficients in  $\mathcal{A}$ . We introduce a map

$$F : [0, 1] \times J^\infty(M) \rightarrow J^\infty(M), \quad (t, u^{i,s}) \mapsto (t^s u^{i,s}),$$

it gives a homotopy from  $s_0 \circ \pi_\infty$  to  $\text{Id}_{J^\infty(M)}$ , where  $s_0$  and  $\pi_\infty$  is the zero section and the structure projection of the bundle  $J^\infty(M)$  respectively, so the induced maps

$$(s_0 \circ \pi_\infty)^*, \quad \text{Id}_{J^\infty(M)}^* : H^*(J^\infty(M), d_{dR}) \rightarrow H^*(J^\infty(M), d_{dR})$$

must be same. But note that when  $d > 0$ , the induced map

$$(s_0 \circ \pi_\infty)^* : H_d^*(J^\infty(M), d_{dR}) \rightarrow H_d^*(J^\infty(M), d_{dR})$$

is zero, so we obtain

$$H^*(\mathcal{S}_{loc}^{>0}, D'_P) \cong H_0^*(J^\infty(M), d_{dR}) \cong H_{dR}^*(M) \cong \mathbb{R}, \quad (3.21)$$

since we have assumed that  $M$  is contractible.

*Proof of Theorem 3.3.6* From Equations (3.21) and (3.20) we obtain

$$H_d^p(\mathcal{S}_{loc}, D'_P) \cong \begin{cases} \wedge^p(\mathbb{R}^n), & d = 0, \\ 0, & d > 0, \end{cases}$$

so the long exact sequence (3.19) implies  $H_{>0}^*(\mathcal{S}) \cong 0$ , then the theorem is proved by applying the long exact sequence (3.18).  $\square$

### 3.5 Bi-Jacobi structures

In this subsection we introduce the notion of compatibility of two Jacobi structures, and use the Lenard-Magri recursion scheme to study integrability of a system which possesses a pair of compatible Jacobi structures, i.e., a bi-Jacobi structure. We also consider properties of a bi-Jacobi structure under reciprocal transformations. Note that in the finite dimensional case, compatible Jacobi structures and their reductions were studied by J. M. Nunes da Costa in [48].

**Definition 3.5.1** *We say that two Jacobi structures  $P, Q \in \mathcal{E}^2$  are compatible if  $[P, Q] = 0$ , and call a compatible pair of Jacobi structures a bi-Jacobi structure. We say that an evolutionary PDE  $X \in \mathcal{E}_{loc}^1$  possesses a bi-Jacobi structure if there exists a bi-Jacobi structure  $(P, Q)$  and two local functionals  $H_0, H_1$  such that*

$$X = [P, H_1] = [Q, H_0]. \quad (3.22)$$

In what follows we assume that both leading terms of  $P$  and  $Q$  are of hydrodynamic type. According to Main Theorem, there exists a reciprocal transformation  $\Phi$  such that  $\Phi(P)$  is local. Then  $\Phi(X)$  is also local, due to the following identity:

$$\Phi(X) = \Phi([P, H_1]) = [\Phi(P), \Phi(H_1)].$$

So without loss of generality we assume that  $P$  is local.

**Lemma 3.5.2** *For a given evolutionary PDE (3.22) that possesses a bi-Jacobi structure, there exist local functionals  $H_2, H_3, \dots \in \mathcal{F}$  such that*

$$[P, H_{k+1}] = [Q, H_k], \quad k = 1, 2, \dots \quad (3.23)$$

and moreover, for any  $k, l \geq 0$  we have

$$P(H_k, H_l) = 0, \quad Q(H_k, H_l) = 0. \quad (3.24)$$

*Proof* By using the Jacobi identity and the second equality of (3.22) we have

$$[P, [Q, H_1]] = -[Q, [P, H_1]] - [H_1, [P, Q]] = -[Q, [P, H_1]] = -[Q, [Q, H_0]] = 0,$$

so from Corollary 3.3.7 it follows the existence of a local functional  $H_2$  such that

$$[P, H_2] = [Q, H_1].$$

By repeating this procedure we can obtain local functionals  $H_2, H_3, \dots$  satisfying (3.23).

The equations (3.24) can be proved by using the recursion relation (3.23) and Magri's trick given in [43].  $\square$

We denote  $X_k = [P, H_k]$ , then  $X_k$ 's are all local, and for any  $k, l \geq 0$  we have

$$[X_k, H_l] = 0, \quad [X_k, X_l] = 0.$$

So in the case when  $\{H_k\}_{k=0}^\infty$  are linearly independent we obtain an integrable hierarchy  $\{X_k\}_{k=1}^\infty$  which has infinitely many common conserved quantities  $\{H_k\}_{k=0}^\infty$ .

The notion of bi-Jacobi structure is a generalization of the one of bihamiltonian structure, the above lemma shows its close relation with integrable systems. An important problem concerning such a relation is the problem of classification of the orbit space of bi-Jacobi structures under the action of Miura type transformations and reciprocal transformations. In the local case, i.e. when both  $P$  and  $Q$  are local, the classification problem has been partially solved in [16, 39]. The main results of these papers are the introduction of the concept of central invariants for any deformation of a semisimple bihamiltonian structures of hydrodynamic type, and proof of the fact that two deformations are equivalent if and only if they possess the same central invariants. In the general nonlocal case the classification problem is far from being solved.

In what follows of this subsection, we proceed to consider properties of a bi-Jacobi structure under reciprocal transformations. To this end we first recall some definitions.

**Definition 3.5.3** *Let  $(P_0, Q_0)$  be a bihamiltonian structure of hydrodynamic type*

$$P_0 \sim \frac{1}{2} g_1^{ij} \theta_i \theta_j^1 + \Gamma_{1,k}^{ij} u^{k,1} \theta_i \theta_j \quad , \quad Q_0 \sim \frac{1}{2} g_2^{ij} \theta_i \theta_j^1 + \Gamma_{2,k}^{ij} u^{k,1} \theta_i \theta_j \quad .$$

*We say that  $(P_0, Q_0)$  is semisimple if the roots of the characteristic equation*

$$\det g_2^{ij} - \lambda g_1^{ij} = 0 \tag{3.25}$$

*are distinct and not constant.*

According to the result of Ferapontov [24], the property of semisimplicity implies that the roots of the characteristic equation (3.25) can be used as local coordinates near every point of  $M$ . We denote them by  $\lambda_1, \dots, \lambda_n$  and call them the canonical coordinates of the bihamiltonian structure  $(P_0, Q_0)$  [16, 39]. The two metrics become diagonal under the canonical coordinates,

$$g_1^{ij}(\lambda) = \delta^{ij} f_i(\lambda), \quad g_2^{ij}(\lambda) = \delta^{ij} \lambda_i f_i(\lambda).$$

From now on in this subsection, we will assume that  $(P_0, Q_0)$  is a semisimple bihamiltonian structure represented in its canonical coordinates.

Let  $(P, Q)$  be a deformation of  $(P_0, Q_0)$ , we introduce the following tensors:

$$A_m^{ij} = \partial_m^j \delta^i(P)_0, \quad B_m^{ij} = \partial_m^j \delta^i(Q)_0, \quad m = 0, 1, 2, \dots \tag{3.26}$$

where  $(W)_0$  denotes the degree zero part of  $W \in \mathcal{A}$ . For example,  $A_0^{ij} = B_0^{ij} = 0$ ,  $A_1^{ij} = g_1^{ij}$ ,  $B_1^{ij} = g_2^{ij}$ , and so on.

**Definition 3.5.4** ([16]) *The central invariants of  $(P, Q)$  are defined by*

$$c_i(\lambda) = \frac{1}{3f_i^2} \left( B_3^{ii} - \lambda_i A_3^{ii} + \prod_{k \neq i} \frac{(B_2^{ki} - \lambda_i A_2^{ki})^2}{f_k(\lambda_k - \lambda_i)} \right), \quad i = 1, \dots, n. \quad (3.27)$$

*It can be shown that  $c_i$  only depends on  $\lambda_i$ .*

We have the following important property of the central invariants:

**Theorem 3.5.5** ([16, 39]) *Let  $(P, Q)$ ,  $(P', Q')$  be two deformations of  $(P_0, Q_0)$ , and  $c_i, c'_i$  ( $i = 1, \dots, n$ ) be their central invariants respectively, then  $(P, Q)$  is equivalent to  $(P', Q')$  via a Miura type transformation if and only if  $c_i = c'_i$  ( $i = 1, \dots, n$ ).*

Let us prove the the main result of this subsection given in Theorem 1.3.

*Proof of Theorem 1.3* Let the bihamiltonian structure  $(P, Q)$  be a deformation of  $(P_0, Q_0)$ , and  $\Phi$  be a reciprocal transformation such that both  $\Phi(P)$  and  $\Phi(Q)$  are local. Since reciprocal transformations induce conformal changes of the flat metrics  $g_1$  and  $g_2$  associated to  $P_0$  and  $Q_0$ , it is easy to see that the canonical coordinates are preserved under such transformations. We denote by  $\tilde{A}_m^{ij}, \tilde{B}_m^{ij}$  the tensors (3.26) associated to  $(\Phi(P), \Phi(Q))$ , then by using the formula (2.39) we can easily obtain

$$\tilde{A}_m^{ij} = \rho^{m+1} A_m^{ij}, \quad \tilde{B}_m^{ij} = \rho^{m+1} B_m^{ij}.$$

By using these relations the theorem follows from the definition of the central invariants immediately.  $\square$

**Remark 3.5.6** *In Theorem 1.3 the condition that  $\Phi(P), \Phi(Q)$  are local is crucial. There are some important bihamiltonian integrable hierarchies which are equivalent via reciprocal transformations but have different central invariants. They are not counter-examples to Theorem 1.3 because these integrable hierarchies possess more than two Jacobi structures, after the reciprocal transformation some of their local Jacobi structures may become nonlocal, and some nonlocal ones may become local. So the resulting hierarchy may still possess a (local) bihamiltonian structure, but this bihamiltonian structure is not the one that is transformed from the original local one, so the theorem cannot apply to such cases. We give an example for this case in the next section, one can find more examples in [1] and [9].*

**Corollary 3.5.7** *Let  $(P, Q)$  be a bihamiltonian structure whose leading term is semi-simple and of hydrodynamic type. For two invertible differential polynomials  $\rho, \rho' \in \mathcal{A}$  satisfying  $\rho \sim \rho'$ , let  $\Phi, \Phi'$  be the reciprocal transformations associated to them respectively. Then if both  $\Phi(P), \Phi(Q)$  are local, there exists a Miura type transformation  $g$  such that*

$$g(\Phi(P)) = \Phi'(P), \quad g(\Phi(Q)) = \Phi'(Q).$$

*Proof* According to Corollary 3.2.6,  $\Phi'(P)$  and  $\Phi'(Q)$  are both local. Suppose  $\rho' = \rho + \partial F$ , then  $\eta = \rho'/\rho = 1 + \tilde{\partial} F$ . Let  $\Psi$  be the reciprocal transformation w.r.t.  $\eta$ , then we have

$$\Psi(\Phi(P)) = \Phi'(P), \quad \Psi(\Phi(Q)) = \Phi'(Q),$$

so  $\Psi$  is a reciprocal transformation converting a local bihamiltonian structure to a local bihamiltonian structure, so the central invariants of  $(\Phi(P), \Phi(Q))$  and  $(\Phi(P'), \Phi(Q'))$  must coincide. On the other hand, it is obvious that the bihamiltonian structures  $(\Phi(P), \Phi(Q))$  and  $(\Phi(P'), \Phi(Q'))$  possess same leading terms, so they are deformations of a same semisimple bihamiltonian structure of hydrodynamic type. Thus according to Theorem 3.5.5 there exists a Miura type transformation  $g$  such that

$$g(\Phi(P)) = \Phi'(P), \quad g(\Phi(Q)) = \Phi'(Q),$$

the corollary is proved. □

We note that the above corollary only ensures the existence of  $g$ , we do not have the explicit form of  $g$  in general.

## 4 Examples

### 4.1 The KdV equation v.s. the Camassa-Holm equation

In this subsection, we investigate the relation between the Jacobi structures of the KdV equation and the Camassa-Holm equation.

The KdV equation

$$u_t = X = 6u u_x - u_{xxx}, \tag{4.1}$$

possesses two compatible local Hamiltonian structures with the following Hamiltonian operators [28, 43, 54]:

$$P_0 = \partial, \quad P_1 = u \partial + \frac{1}{2} u_x - \frac{1}{4} \partial^3.$$

Denote  $R = P_1 \circ P_0^{-1}$ , and  $P_k = R^k \circ P_0$ , then  $R$  is a hereditary strong symmetry [27], and all  $P_k$ 's are Hamiltonian operators which are, however, nonlocal for  $k \geq 2$ . Among these nonlocal Hamiltonian operators, there are two operators that correspond

to Jacobi structures:

$$\begin{aligned}
P_2 &= u^2 \partial + u u_x - \frac{1}{4} u_x \partial^{-1} u_x \\
&\quad - \frac{1}{2} u \partial^3 + \frac{3}{4} u_x \partial^2 + \frac{1}{2} u_{xx} \partial + \frac{1}{8} u_{xxx} \quad + \frac{1}{16} \partial^5, \\
P_3 &= u^3 \partial + \frac{3}{2} u^2 u_x \\
&\quad - \frac{1}{16} (6u u_x - u_{xxx}) \partial^{-1} u_x - \frac{1}{16} u_x \partial^{-1} (6u u_x - u_{xxx}) \\
&\quad - \frac{3}{4} u^2 \partial^3 + \frac{9}{4} u u_x \partial^2 + \frac{3}{2} u u_{xx} + \frac{17}{16} u_x^2 \partial \\
&\quad + \frac{3}{8} u u_{xx} + \frac{11}{16} u_x u_{xx} \quad + \frac{3}{16} u \partial^5 + \frac{15}{32} u_x \partial^4 \\
&\quad + \frac{5}{8} u_{xx} \partial^3 + \frac{15}{32} u_{xxx} \partial^2 + \frac{3}{16} u_{4x} \partial + \frac{1}{32} u_{5x} \quad - \frac{1}{64} \partial^7
\end{aligned}$$

We denote by  $Q_i$  ( $i = 0, 1, 2, 3$ ) the quasi-local bivectors corresponding to  $P_i$  ( $i = 0, 1, 2, 3$ )

$$Q_i = \frac{1}{2} \int \theta P_i(\theta) dx,$$

then they read

$$\begin{aligned}
Q_0 &= \frac{1}{2} \int \theta \theta' dx, \\
Q_1 &= \frac{1}{2} \int u \theta \theta' + \frac{3}{2} \theta \theta''' dx, \\
Q_2 &= \frac{1}{2} \int u^2 \theta \theta' + \frac{1}{4} u_x \theta \zeta + \dots dx, \\
Q_3 &= \frac{1}{2} \int u^3 \theta \theta' + \frac{3}{4} u u_x \theta \zeta + \dots dx,
\end{aligned}$$

where we omitted some terms with higher degrees. Here an interesting problem arises: does there exist more Jacobi structures other than  $Q_0, Q_1, Q_2, Q_3$  for the KdV equation (4.1)? We conjecture that the answer is no, and we will investigate this problem in a separate publication [41].

Let  $\rho \in \mathcal{A}$  be an invertible differential polynomial, and  $\Phi$  be the reciprocal transformation w.r.t.  $\rho$ . It is easy to see that to ensure the locality of  $\Phi(X)$  the differential polynomial  $\rho$  must be a conserved density of the KdV hierarchy. From the quasi-triviality of the KdV hierarchy [40] it follows that the conserved densities are uniquely determined by their leading terms (up to a total differential). We assume that the leading term of  $\rho$  is  $u^\kappa$  ( $\kappa \in \mathbb{R}$ ), then by using Lemma 3.1.7 and the results of Section 3.2 one can obtain the following assertions:

- (a)  $\Phi(Q_0)$  is local  $\Leftrightarrow \kappa = 0, 2$ ;    (b)  $\Phi(Q_1)$  is local  $\Leftrightarrow \kappa = 0, 1$ ;  
(c)  $\Phi(Q_2)$  is local  $\Leftrightarrow \kappa = \frac{1}{2}, -\frac{1}{2}$ ;    (d)  $\Phi(Q_3)$  is local  $\Leftrightarrow \kappa = \frac{1}{2}, -\frac{3}{2}$ .

The case  $\kappa = 0$  is trivial. When  $\kappa = 2$ , we can choose  $\rho = u^2$ , the transformed equation  $\Phi(X)$  is analyzed in [40] and [42]. It is shown in [42] that  $\Phi(X)$  has only one local Hamiltonian structure. Now we know that it also has (at least) three Jacobi structures. The case when  $\kappa = 1$  is similar.

We are interested in the case when  $\kappa = \frac{1}{2}$ . For this we first need to find out the full expression of  $\rho$ . It is well known that the following Miura transformation

$$u = v^2 + v_x$$

converts the KdV equation (4.1) to the modified KdV equation

$$v_t = 6v^2 v_x - v_{xxx} = 2v^3 - v_{xx} \quad ,$$

so we can choose the conserved density  $\rho = 2v$ . Here the factor 2 is not essential, we add it just for the convenience of relating the KdV equation with the standard Camassa-Holm equation. After fixing this  $\rho$ , one obtains  $\Phi(Q_i)$  ( $i = 0, 1, 2, 3$ ) with  $\Phi(Q_2)$  and  $\Phi(Q_3)$  being local.

**Theorem 4.1.1** *There exists a Miura type transformation  $g$  such that*

$$(g(\Phi(Q_2)), g(\Phi(Q_3)))$$

*gives the bihamiltonian structure of the Camassa-Holm hierarchy [7, 8, 26, 27].*

*Proof* We recall some facts about the Camassa-Holm (CH) equation first. The CH equation

$$m_s = 2m w_y + m_y w, \quad m = w - w_{yy} \quad (4.2)$$

has two local Hamiltonian structures

$$J_0 = \partial_y - \partial_y^3, \quad J_1 = m \partial_y + \frac{1}{2} m_y$$

and infinitely many nonlocal Hamiltonian structures  $J_k = J_1 J_0^{-1} J_0$  among which  $J_2$  and  $J_3$  correspond to Jacobi structures. The CH hierarchy can be obtained by using the recursion operator  $J_1 J_0^{-1}$ .

The simplest way to describe the CH hierarchy is to consider the isospectral problem

$$\phi_{yy} = \frac{1}{4} - m \lambda \quad \phi, \quad \phi_s = -B(\lambda)\phi_y + \frac{1}{2} B_y(\lambda)\phi, \quad (4.3)$$

where  $B(\lambda)$  is a Laurent polynomial of the spectral parameter  $\lambda$ . For example, when

$$B(\lambda) = w + \frac{1}{2\lambda},$$

we obtain the CH equation (4.2) from the compatibility condition  $\phi_{yys} = \phi_{syy}$ .

The CH hierarchy has a conserved density  $\tilde{\rho} = \sqrt{m}$ , we denote the associated reciprocal transformation by  $\tilde{\Phi}$ . Let  $\psi = m^{\frac{1}{4}} \phi$ , then the  $y$ -part of the isospectral problem (4.3) becomes

$$\psi_{xx} = (u - \lambda) \psi, \quad (4.4)$$

where  $x = \tilde{y}$  is defined by

$$d\tilde{y} = \tilde{\rho} dy + w\sqrt{m} ds, \quad d\tilde{s} = ds$$

and the function  $u$  reads

$$u = \frac{1}{4m} + \frac{m_{xx}}{4m} - \frac{3m_x^2}{16m^2}. \quad (4.5)$$

Note that the equation (4.4) is exactly the  $x$ -part of the isospectral problem of the KdV equation (4.1). In fact, if we take the  $t$ -part as

$$\psi_t = (4\lambda + 2u)\psi_x - u_x\psi,$$

then the compatibility condition  $\psi_{xxt} = \psi_{txx}$  gives us the KdV equation (4.1). So we see that the reciprocal transformation  $\tilde{\Phi}$  together with the Miura type transformation (4.5) transforms the CH hierarchy to the negative flows of the KdV hierarchy. Let  $\Phi'$  be the inverse of  $\tilde{\Phi}$ , then the reciprocal transformation  $\Phi'$  is defined by

$$\rho' = \tilde{\rho}^{-1} = 2\sqrt{u} + \dots$$

Note that  $\rho'$  must be a conserved density of the KdV hierarchy, so by using the quasi-triviality property of the KdV hierarchy [40] we have  $\rho' \sim \rho$ .

We denote the quasi-local bivectors corresponding to  $J_i$  ( $i = 0, 1, 2, 3$ ) by  $K_i$  ( $i = 0, 1, 2, 3$ ). Since  $\rho'$  is a conserved density with leading term  $2\sqrt{u}$ , we know from Corollary 3.2.6 that the Jacobi structures  $\Phi'(Q_2), \Phi'(Q_3)$  are also local. By a straightforward computation, one can show that the leading terms of  $\Phi'(Q_2), \Phi'(Q_3)$  coincide with that of  $K_1, K_0$  (after the Miura type transformation (4.5)). On the other hand, the main results of [42] imply that the Hamiltonian structures of the CH equation are uniquely determined by their leading terms, so we have

$$\Phi'(Q_2) = K_1, \quad \Phi'(Q_3) = K_0.$$

A similar argument shows that

$$\tilde{\Phi}(K_2) = Q_1, \quad \tilde{\Phi}(K_3) = Q_0,$$

thus we have  $\Phi'(Q_0) = K_3, \Phi'(Q_1) = K_2$ .

Finally, due to Corollary 3.5.7 there exists a Miura type transformation  $g$  such that

$$g(\Phi(Q_2)) = \Phi'(Q_2), \quad g(\Phi(Q_3)) = \Phi'(Q_3),$$

the theorem is proved.  $\square$

The relation between the Jacobi structures of the KdV equation and that of the CH equation has a multi-component generalization, which relates the Jacobi structures of the  $r$ -KdV-CH hierarchy associated to the parameters  $(a_0, \dots, a_r) \in (\mathbb{R}^{r+1})^\times$  with the Jacobi structures of the  $r$ -KdV-CH hierarchy associated to the parameters  $(a_r, \dots, a_0)$ , see [9] for more details.

In [1], Abenda and Grava investigated the relations of the Whitham equations for the CH equation with that of the KdV equation. They found similar relations between the local and non-local Hamiltonian structures of these Whitham equations.

## 4.2 Bihamiltonian structures associated to Frobenius manifolds

In this subsection we consider reciprocal transformation of the bihamiltonian structure defined on the jet space of a Frobenius manifold. The notion of Frobenius manifold was introduced by Dubrovin[14, 15] as a geometrical interpretation of the WDVV equations that arose in the study of 2D topological field theory [13, 51]. On any Frobenius manifold  $M$  there is defined a flat metric  $\eta$ , and on each of its tangent spaces there is a Frobenius algebra structure which depends smoothly on the point of the manifold. The structure constants of the Frobenius algebras can be represented by a function  $F$ , called the potential of the Frobenius manifold, of the flat coordinates  $v^1, \dots, v^n$  of  $M$ . These flat coordinates can be chosen in such a way that the vector field  $e_1 = \frac{\partial}{\partial v^1}$  gives the unity element of the Frobenius algebras. Denote by  $\eta_{ij}$  the components of the flat metric in the flat coordinates, and  $(\eta^{ij}) = (\eta_{ij})^{-1}$ , then the operation of multiplication of the Frobenius algebras on the tangent spaces is given by

$$\frac{\partial}{\partial v^i} \cdot \frac{\partial}{\partial v^j} = c_{ij}^k \frac{\partial}{\partial v^k}, \quad c_{ij}^k = \eta^{kl} \frac{\partial^3 F(v)}{\partial v^l \partial v^i \partial v^j}.$$

Note that we have  $c_{1j}^k = \delta_j^k$ . The associativity condition of the Frobenius algebras yields a system of nonlinear PDEs for the potential  $F$ . This function also satisfies certain quasi-homogeneity condition which is represented by an Euler vector field  $E$  of the form

$$E = \sum_{i=1}^n E^i \frac{\partial}{\partial v^i} = \sum_{i=1}^n (d_i v^i + r_i) \frac{\partial}{\partial v^i}, \quad \text{with } d_1 = 1 \quad (4.6)$$

satisfying

$$\partial_E F = (3 - d)F + \frac{1}{2} A_{ij} v^i v^j + B_i v^i + C. \quad (4.7)$$

Here  $d_i, r_i, d, A_{ij}, B_i, C$  are some constants,  $A_{1i} = A_{i1} = \prod_{l=1}^n \eta_{il} r_l$ , and the constant  $d$  is called the charge of the Frobenius manifold. The Euler vector field is so normalized that  $r_k = 0$  when  $d_k \neq 0$ .

In what follows of this subsection we assume that the flat coordinates can be normalized so that the metric  $\eta$  takes the form

$$\eta_{ij} = \delta_{i, n+1-j}, \quad (4.8)$$

and the potential  $F(v)$  has the expression

$$F(v) = \frac{1}{2}(v^1)^2 v^n + \frac{1}{2} \sum_{k,l=2}^n \eta_{kl} v^1 v^k v^l + f(v^2, \dots, v^n).$$

Then the constants  $d_i$  satisfy the following duality property:

$$d_i + d_{n+1-i} = 2 - d.$$

We also assume that

$$r_1 = \dots = r_n = 0,$$

which implies  $A_{1i} = 0$  ( $i = 1, \dots, n$ ) and  $B_1 = 0$ .

The flat metric  $\eta$  yields a Hamiltonian structure of hydrodynamic type on the jet space of  $M$  given by the Hamiltonian operator

$$P_1^{ij} = \eta^{ij} \partial_x, \quad i, j = 1, \dots, n. \quad (4.9)$$

There is another compatible Hamiltonian structure which is associated to the intersection form  $(g^{ij})$  of the Frobenius manifold defined by

$$g^{ij} = E^k c_k^{ij}, \quad c_k^{ij} = \eta^{il} c_{lk}^j. \quad (4.10)$$

The Hamiltonian operator has the form

$$P_2^{ij} = g^{ij}(v) \partial_x + \Gamma_k^{ij}(v) v_x^k, \quad (4.11)$$

where  $\Gamma_k^{ij} = -g^{il} \Gamma_{lk}^j$  are the contravariant components of the Levi-Civita connection of the metric  $(g_{ij}) = (g^{ij})^{-1}$  defined on the open subset of  $M$  where  $\det(g^{ij}(v)) \neq 0$ .

Let us consider the reciprocal transformation  $\Phi$  w.r.t. the function  $\rho = v^n$  of the bihamiltonian structure  $(P_1, P_2)$ . By using Lemma 3.1.7 we know that the bihamiltonian structure is transformed to a bi-Jacobi structure in general. However in this particular case the bihamiltonian structure is transformed to a local bi-Jacobi structure, i.e. again a bihamiltonian structure. In the coordinates  $v^1, \dots, v^n$  this bihamiltonian structure is given by

$$\Phi(P_1)^{ij} = \tilde{\eta}^{ij} \partial_{\tilde{x}} + \tilde{\Gamma}_{1,k}^{ij} v_{\tilde{x}}^k, \quad \Phi(P_2)^{ij} = \tilde{g}^{ij} \partial_{\tilde{x}} + \tilde{\Gamma}_{k,2}^{ij} v_{\tilde{x}}^k, \quad (4.12)$$

where  $\tilde{\eta}^{ij} = \rho^2 \eta^{ij}$ ,  $\tilde{g}^{ij} = \rho^2 g^{ij}$ , and  $\tilde{\Gamma}_{1,k}^{ij}, \tilde{\Gamma}_{2,k}^{ij}$  are the contravariant components of the Levi-Civita connection of  $(\tilde{\eta}_{ij}), (\tilde{g}_{ij})$  respectively.

Note that the metrics  $\tilde{\eta}$  and  $\tilde{g}$  are also flat. It is easy to see that the first metric  $\tilde{\eta}$  has the following system of flat coordinates:

$$\tilde{v}^1 = \frac{1}{2} \frac{\eta_{kl} v^k v^l}{v^n}, \quad \tilde{v}^i = \frac{v^i}{v^n} \quad (i \neq 1, n), \quad \tilde{v}^n = -\frac{1}{v^n}. \quad (4.13)$$

In these flat coordinates the metric  $\tilde{\eta}$  has the contravariant components

$$\tilde{\eta}^{ij}(\tilde{v}) = \eta^{ij}. \quad (4.14)$$

It is noticed in [52] that the metrics  $\tilde{\eta}(\tilde{v})$  and  $\tilde{g}(\tilde{v})$  coincide with the flat metric and the intersection form of the Frobenius manifold  $\tilde{M}$  with the potential

$$\tilde{F}(\tilde{v}) = (\tilde{v}^n)^2 F(v) + \frac{1}{2} \eta_{kl} \tilde{v}^1 \tilde{v}^k \tilde{v}^l. \quad (4.15)$$

Here in the r.h.s. of the above formula the variables  $v^1, \dots, v^n$  of  $F(v)$  are understood to be dependent on  $\tilde{v}^1, \dots, \tilde{v}^n$  via (4.13). The Euler vector field  $\tilde{E} = \sum_{i=1}^n \tilde{E}^i \frac{\partial}{\partial \tilde{v}^i}$  of the Frobenius manifold  $\tilde{M}$  coincides with  $E$ , i.e.

$$\tilde{E} = \sum_{i=1}^n \tilde{E}^i \frac{\partial}{\partial \tilde{v}^i} = \sum_{i=1}^n E^i \frac{\partial}{\partial v^i}.$$

and the charge of  $\tilde{M}$  is  $\tilde{d} = 2 - d$ .

The construction of the Frobenius manifold  $\tilde{M}$  from  $M$  by the above formula (4.13)–(4.15) was given by Dubrovin in [15], it corresponds to a special symmetry of the WDVV equations called the inversion symmetry. The above fact gives a natural interpretation of this symmetry of the WDVV equations in terms of a special reciprocal transformation of the associated bihamiltonian structure on the jet space of the Frobenius manifold. Note that under certain additional conditions a bihamiltonian structure of hydrodynamic type determines a Frobenius manifold [20]. In [52] the transformation rule of the associated bihamiltonian integrable hierarchy, called the principal hierarchy of the Frobenius manifold, under the above reciprocal transformation is given, it shows that the reciprocal transformation induces a Legendre transformation of the tau function of the principal hierarchy.

## 5 Conclusion

In the preceding sections we introduce the notion of infinite dimensional Jacobi structures which provides a geometrical characterization of an important class of nonlocal Hamiltonian structures. The invariance of the space of Jacobi structures under reciprocal transformations enables one to study normal forms of Jacobi structures under Miura type transformations and reciprocal transformations, and to discover possible hidden relations between some important evolutionary PDEs via the study of their local and nonlocal Hamiltonian structures. For a Jacobi structure with leading term of hydrodynamic type, we obtain its normal form by computing the associated Lichnerowicz-Jacobi cohomologies.

From the point view of integrable evolutionary PDEs, our next important task is to consider the problem of classification of bi-Jacobi structures with leading terms of hydrodynamic type. As we have shown in Section 3.5, under certain appropriate conditions an evolutionary PDE which has a bi-Jacobi structure possesses an infinite number of commuting flows and conservation laws. Thus bi-Jacobi structures just like bihamiltonian structures are closely related to integrable evolutionary PDEs. To

classify bi-Jacobi structures of hydrodynamic type, one must classify first their leading terms, i.e. pairs of compatible conformally flat metrics, which is a pure differential geometric problem; then the classification problem becomes a deformation problem, which is controlled by some *bi-Jacobi cohomologies*. The computation of these bi-Jacobi cohomologies is highly nontrivial, new techniques are expected.

The reciprocal transformations that we consider in this paper only change the spatial variable  $x$  of a system of evolutionary PDEs  $u_t^i = K^i(u, u_x, \dots)$ ,  $i = 1, \dots, n$ . It is also important to consider more general class of reciprocal transformations of the form

$$d\tilde{x} = \rho_1 dx + \sigma_1 dt, \quad d\tilde{t} = \rho_2 dx + \sigma_2 dt, \quad (5.1)$$

they change both the time and spatial variables. Here  $\frac{\partial \rho_i}{\partial t} = \frac{\partial \sigma_i}{\partial x}$ ,  $i = 1, 2$  are two conservation laws of the evolutionary PDEs which satisfy certain non-degeneracy conditions. Such reciprocal transformations can be represented as the composition of two transformations of the form (1.6) and one of the form

$$d\tilde{x} = dt, \quad d\tilde{t} = dx, \quad (5.2)$$

i.e. a transformation that exchanges the role of the time and spatial variables of the evolutionary PDEs. The transformation rule of the Hamiltonian structures of a system of Hamiltonian evolutionary PDEs under the reciprocal transformation (5.2) was considered in [50], where it was shown that for Hamiltonian systems of hydrodynamic type the reciprocal transformation (5.2) preserves the forms of the systems and their Hamiltonian structures. It was further shown in [53] that the reciprocal transformation (5.2) also preserves the bihamiltonian structures of bihamiltonian systems of hydrodynamic type. We conjecture that for a general Hamiltonian (resp. bihamiltonian) system with dispersion terms the reciprocal transformation (5.2) preserves the locality property of the Hamiltonian and bihamiltonian structures. If this conjecture holds true, we could consider the problem of classification of bihamiltonian structures and the associated hierarchies of integrable evolutionary PDEs under Miura type transformations and also reciprocal transformations which preserve the locality property. It is expected that the moduli spaces that arise in such a classification scheme are much simple than that of the bihamiltonian integrable hierarchies under the action of Miura type transformations. A preliminary study suggests that every hierarchy of scalar evolutionary PDEs with leading terms of hydrodynamic type and a bihamiltonian structure is equivalent to the usual KdV hierarchy under Miura type transformations and reciprocal transformations of the form (5.1). We will discuss this issue in separate publications.

The action of reciprocal transformation (5.2) on general Jacobi structures is still an open problem. The results of [25] suggest that reciprocal transformations of the form (5.2) may convert Jacobi structures to more general nonlocal Hamiltonian structures which are not rigorously defined yet. In a recent paper [33], Kersten *et al* sketch out a new geometric framework to construct Hamiltonian operators for generic, non-evolutionary PDEs, we hope that their approach will be helpful to the study of this problem.

The notion of Jacobi structures defined in the present paper is only for  $1 + 1$  dimensional evolutionary PDEs. It is not hard to generalize it to higher spatial dimensional cases. To do so one only needs to replace the jet space  $J^\infty(M)$  used in Section 2 by more general jet spaces. Some results of the present paper still hold true for Jacobi structures in higher spatial dimensional cases, while some results do not. There are many interesting open problems in this aspect.

**Acknowledgments.** The authors thank Boris Dubrovin for his interests in and advises on the work of the paper, and the referee for helpful comments and suggestions. Part of the work was done in SISSA during the authors' visit to it, they thank its hospitality and financial support. The work is partially supported by the National Basic Research Program of China (973 Program) No. 2007CB814800, the NSFC No.10631050, 10801084 and 11071135.

## References

- [1] S. Abenda, T. Grava, Modulation of Camassa-Holm equation and reciprocal transformations, *Ann. Inst. Fourier (Grenoble)* 55 (2005) 1803–1834.
- [2] S. Abenda, T. Grava, Reciprocal transformations and flat metrics on Hurwitz spaces, *J. Phys. A* 40 (2007) 10769–10790.
- [3] S. Abenda, Reciprocal transformations and local Hamiltonian structures of hydrodynamic-type systems, *J. Phys. A* 42 (2009) 095208.
- [4] V. I. Arnold, B. A. Khesin, *Topological methods in hydrodynamics*, Applied Mathematical Sciences 125, Springer-Verlag, New York, 1998.
- [5] A. Barakat, On the moduli space of deformations of bihamiltonian hierarchies of hydrodynamic type, *Adv. Math.* 219 (2008) 604–632.
- [6] K. H. Bhaskara, K. Viswanath, *Poisson algebras and Poisson manifolds*, Pitman Research Notes in Mathematics Series, 174. Longman Scientific & Technical, Harlow; copublished in the United States with John Wiley & Sons, Inc., New York, 1988.
- [7] R. Camassa, D. D. Holm, An integrable shallow water equation with peaked solitons, *Phys. Rev. Lett.* 71 (1993) 1661–1664.
- [8] R. Camassa, D. D. Holm, J. M. Hyman, A new integrable shallow water equation, *Adv. Appl. Mech.* 31 (1994) 1–33.
- [9] M. Chen, S.-Q. Liu, Y. Zhang, Hamiltonian structures and their reciprocal transformations for the  $r$ -KdV-CH hierarchy, *J. Geom. Phys.* 59 (2009) 1227–1243.

- [10] P. Dazord, A. Lichnerowicz, C. Marle, Structure locale des variétés de Jacobi, *J. Math. Pures Appl.* (9) 70 (1991) 101–152.
- [11] L. Degiovanni, F. Magri, V. Sciacca, On deformation of Poisson manifolds of hydrodynamic type, *Comm. Math. Phys.* 253 (2005) 1–24.
- [12] L. A. Dickey, Soliton equations and Hamiltonian systems, Second edition, *Advanced Series in Mathematical Physics* 26, World Scientific Publishing Co. Inc., River Edge, NJ, 2003.
- [13] R. Dijkgraaf, E. Verlinde, H. Verlinde, Topological strings in  $d < 1$ , *Nucl. Phys. B* 352 (1991) 59–86.
- [14] B. Dubrovin, Integrable systems and classification of 2-dimensional topological field theories, In *Integrable systems*, Proceedings of Luminy 1991 conference dedicated to the memory of J.-L. Verdier, eds. O. Babelon, O. Cartier, Y. Kosmann-Schwarbach, Birkhäuser, 1993.
- [15] B. Dubrovin, Geometry of 2D topological field theories, Integrable systems and quantum groups (Montecatini Terme, 1993) 120–348, *Lecture Notes in Math.* 1620, Springer, Berlin, 1996.
- [16] B. Dubrovin, S.-Q. Liu, Y. Zhang, On Hamiltonian perturbations of hyperbolic systems of conservation laws. I. Quasi-triviality of bi-Hamiltonian perturbations, *Comm. Pure Appl. Math.* 59 (2006) 559–615.
- [17] B. Dubrovin, S.-Q. Liu, Y. Zhang, Frobenius manifolds and central invariants for the Drinfeld-Sokolov bi-Hamiltonian structures, *Adv. Math.* 219 (2008) 780–837.
- [18] B. A. Dubrovin, S. P. Novikov, Hamiltonian formalism of one-dimensional systems of the hydrodynamic type and the Bogolyubov-Whitham averaging method, *Dokl. Akad. Nauk SSSR* 270 (1983) 781–785.
- [19] B. A. Dubrovin, S. P. Novikov, Poisson brackets of hydrodynamic type, *Dokl. Akad. Nauk SSSR* 279 (1984) 294–297.
- [20] B. Dubrovin, Flat pencils of metrics and Frobenius manifolds, In: *Proceedings of 1997 Taniguchi Symposium “Integrable Systems and Algebraic Geometry”*, Editors M.-H. Saito, Y. Shimizu and K. Ueno, 47–72. World Scientific, 1998.
- [21] B. Dubrovin, Y. Zhang, Normal forms of integrable PDEs, Frobenius manifolds and Gromov-Witten invariants, arXiv: math/0108160.
- [22] L. D. Faddeev, L. A. Takhtajan, Hamiltonian methods in the theory of solitons, Translated from the Russian by A. G. Reyman, *Springer Series in Soviet Mathematics*, Springer-Verlag, Berlin, 1987.

- [23] E. V. Ferapontov, Conformally flat metrics, systems of hydrodynamic type and nonlocal Hamiltonian operators, *Uspekhi Mat. Nauk* 50 (1995), 175–176, translation in *Russian Math. Surveys* 50 (1995) 811–813.
- [24] E. V. Ferapontov, Compatible Poisson brackets of hydrodynamic type, *Kowalevski Workshop on Mathematical Methods of Regular Dynamics (Leeds, 2000)*, *J. Phys. A* 34 (2001) 2377–2388.
- [25] E. V. Ferapontov, M. V. Pavlov, Reciprocal transformations of Hamiltonian operators of hydrodynamic type: nonlocal Hamiltonian formalism for linearly degenerate systems, *J. Mathematical Phys.* 44 (2003) 1150–1172.
- [26] B. Fuchssteiner, Some tricks from the symmetry-toolbox for nonlinear equations: generalizations of the Camassa-Holm equation, *Phys. D* 95 (1996) 229–243.
- [27] B. Fuchssteiner, A. S. Fokas, Symplectic structures, their Bäcklund transformations and hereditary symmetries, *Phys. D* 4 (1981/82) 47–66.
- [28] C. S. Gardner, Korteweg-de Vries equation and generalizations, IV, The Korteweg-de Vries equation as a Hamiltonian system, *J. Mathematical Phys.* 12 (1971) 1548–1551.
- [29] E. Getzler, A Darboux theorem for Hamiltonian operators in the formal calculus of variations, *Duke Math. J.* 111 (2002) 535–560.
- [30] F. Guedira, A. Lichnerowicz, Géométrie des algèbres de Lie locales de Kirillov, *J. Math. Pures Appl.* (9) 63 (1984) 407–484.
- [31] D. Iglesias, J. C. Marrero, Generalized Lie bialgebroids and Jacobi structures, *J. Geom. Phys.* 40 (2001) 176–199.
- [32] P. Kersten, I. Krasil’shchik, A. Verbovetsky, Hamiltonian operators and  $\ell^*$ -coverings, *J. Geom. Phys.* 50 (2004) 273–302.
- [33] P. Kersten, I. Krasil’shchik, A. Verbovetsky, R. Vitolo, Hamiltonian structures for general PDEs, arXiv: 0812.4895.
- [34] A. A. Kirillov, Local Lie algebras, *Uspekhi Mat. Nauk* 31 (1976) 57–76.
- [35] M. D. Kruskal, R. M. Miura, C. S. Gardner, N. J. Zabusky, Korteweg-de Vries equation and generalizations, V, Uniqueness and nonexistence of polynomial conservation laws, *J. Mathematical Phys.* 11 (1970) 952–960.
- [36] M. de León, B. López, J. C. Marrero, E. Padrón, On the computation of the Lichnerowicz-Jacobi cohomology, *J. Geom. Phys.* 44 (2003) 507–522.
- [37] A. Lichnerowicz, Les variétés de Poisson et leurs algèbres de Lie associées, *J. Differential Geometry* 12 (1977) 253–300.

- [38] A. Lichnerowicz, Les variétés de Jacobi et leurs algèbres de Lie associées, *J. Math. Pures Appl.* (9) 57 (1978) 453–488.
- [39] S.-Q. Liu, Y. Zhang, Deformations of semisimple bihamiltonian structures of hydrodynamic type, *J. Geom. Phys.* 54 (2005) 427–453.
- [40] S.-Q. Liu, Y. Zhang, On quasi-triviality and integrability of a class of scalar evolutionary PDEs, *J. Geom. Phys.* 57 (2006) 101–119.
- [41] S.-Q. Liu, Y. Zhang, On quasi-triviality of Jacobi structures of a class of scalar evolutionary PDEs, in preparation.
- [42] S.-Q. Liu, C.-Z. Wu, Y. Zhang, On properties of Hamiltonian structures for a class of evolutionary PDEs, *Lett. Math. Phys.* 84 (2008) 47–63.
- [43] F. Magri, A simple model of the integrable Hamiltonian equation, *J. Mathematical Phys.* 19 (1978) 1156–1162.
- [44] A. Ya. Maltsev, S. P. Novikov, On the local systems Hamiltonian in the weakly nonlocal Poisson brackets, *Phys. D* 156 (2001) 53–80.
- [45] R. M. Miura, Korteweg-de Vries equation and generalizations, I, A remarkable explicit nonlinear transformation, *J. Mathematical Phys.* 9 (1968) 1202–1204.
- [46] A. Nijenhuis, R. Richardson, Cohomology and deformations in graded Lie algebras, *Bull. Amer. Math. Soc.* 72 (1966) 1–29.
- [47] A. Nijenhuis, R. Richardson, Deformation of Lie algebra structures, *J. Math. Mech.* 17 (1967) 89–105.
- [48] J. M. Nunes da Costa, Compatible Jacobi manifolds: geometry and reduction, *J. Phys. A* 31 (1998) 1025–1033.
- [49] P. J. Olver, Applications of Lie groups to differential equations, Second edition, Graduate Texts in Mathematics 107, Springer-Verlag, New York, 1993.
- [50] M. V. Pavlov, Conservation of the “forms” of Hamiltonian structures upon linear substitution for independent variables, *Math. Notes* 57 (1995) 489–495
- [51] E. Witten, On the structure of the topological phase of two-dimensional gravity, *Nucl. Phys. B* 340 (1990) 281–332.
- [52] D. Xu, Y. Zhang, On symmetries of the WDVV equations, preprint arXiv: 1002.0034.
- [53] T. Xue, Y. Zhang, Bihamiltonian systems of hydrodynamic type and reciprocal transformations, *Lett. Math. Phys.* 75 (2006) 79–92.
- [54] V. E. Zaharov, L. D. Faddeev, The Korteweg-de Vries equation is a fully integrable Hamiltonian system, *Funkcional. Anal. i Priložen.* 5 (1971) 18–27.